First record of the Brassy Chub *Kyphosus vaigiensis* (Pisces: Kyphosidae) from the Mediterranean coast of Libya

Amani FITORI1, Ali EL-FITURI2 and Daniel GOLANI3*

¹Department of Marine Resource, Faculty of Natural Resource, Tobruk University, Libya ² Marine Biology Research Center, Tripoli, Libya.

³Department of Ecology, Evolution and Behavior, The Hebrew University of Jerusalem, Israel

*Corresponding author: dani.golani@mail.huji.ac.il

In the present paper we report the first record of the Brassy Chub Kyphosus vaigiensis (Quoy and Giamard, 1825) from the Mediterranean coast of Libya. It is assumed that this species arrived into the Mediterranean from the Red Sea via the Suez Canal. We stress the need of taxonomic study on the genus Kyphosus in the Mediterranean.

Key words: Kyphosus vaigiensis; Mediterranean; Libya; invasive

INTRODUCTION

The Mediterranean Sea, being an enclosed sea located between Europe and North Africa, has become a hot spot for non-native fish species. A recent publication (GOLANI *et al.*, 2021) enumerated 188 non-indigenous fish species in the Mediterranean; this is an addition of 107 species to the list published less than two decades ago (GOLANI *et al.*, 2002).

New fish enter the Mediterranean in three manners: a) Atlantic species expand their distribution via Gibraltar, b) Red Sea species cross the Suez Canal (Lessepsian migrants) and c) species reach the Mediterranean via direct human activity such as aquaculture and aquarium escapees.

In the present paper, we report the first record of the Brassy Chub *Kyphosus vaigiensis* (Quoy and Giamard, 1825) from the coast of Libya. This record contributes additional evidence of the spread of this species in the Mediterranean.

MATERIAL AND METHODS

On August 2021 two specimens were collected by trammel net at ca. 20 m at location 32°42′17.32″N 21°01′10.20″E (Fig. 1). The specimens were photographed (Fig. 2) but were not saved. All counts and measurements follow HUBBS & LAGLER (1947).




Fig. 1 Map of the site, indicated by a red star, where Kyphosus vaigiensis was captured (Talamitha region, Libya).

RESULTS

Short description (Based on captured specimen – Fig. 2)

Body oblong, moderately deep, its depth 40.1% in SL, small head (21.4% in SL). Small mouth, its jaws terminate at the vertical of interior of eye. Snout length slightly larger than eye diameter. Dorsal fin with IX-X spines and 13-14 rays; Anal fin with III spines and 12-13 rays; Pectoral fin short with 18-19 rays. Pelvic fin origin behind pectoral fin origin with I spine and 5 rays. Caudal fin forked. Color: Body brownishgrey. Back dark, becoming lighter on the ventral surface. Thin horizontal lines on the flank. The posterior margin of the operculum and all fins slightly darker than rest of the body.



Fig. 2 Kyphosus vaigiensis collected from the coast of Libya

DISCUSSION

The family of Kyphosidae, Sea Chubs, consists of 17 species in three genera. According to NELSON *et al.* (2016) the genus *Kyphosus* consists of 15 species, while KNUDSEN & CLEMENTS (2013) enumerated only eleven species in this genus. The morphology of most of the *Kyphosus* species is quite similar, thus having a complicated taxonomy with numerous synonyms and

misidentifications. As a result, several names of the genus Kyphosus appear in literature as occurring in the Mediterranean, such as K. bosquii (LACEPÉDE, 1802) (SAKAI, 2016; ORSI-RELINI, 2017) which according to KNUDSEN & CLEMENTS (2013) is a synonym of the Indo-Pacific K. biggibus Lacepéde, 1801 and the Mediterranean records are most likely misidentifications. TOR-TONESE (1986) and FRANCOUR & MOUINE (2008) reported K. spectator (Linnaeus, 1766) and later LIGAS et al., (2011) and KIPARISSIS et al., (2012) reported K. saltatrix (Linnaeus, 1758); both are synonyms of K. sectatrix (Linnaeus, 1758) (KNUDSEN & CLEMENTS, 2013). Another Atlantic species, K. incisor (Cuvier, 1831), was recorded from the Mediterranean by ORSI-RELINI (2017), but according to KNUDSEN & CLEMENTS (2013), it is a synonym of *K. viagiensis*.

The species K. vaigiensis has a world-wide distribution and it has been reported throughout the Mediterranean (AL MABRUK et al., 2021). The origin of K. vaigiensis in the Mediterranean is not clear. It was first recorded from the Mediterranean from Almuñécar, Granada, Spain as K. incisor in 1998 (AZZURRO et al., 2013); thus suggesting a range expansion from the Atlantic Ocean. However, most of the specimens have been recorded from the Levant, including Israel (GOREN et al., 2016) and central Mediterranean as K. vaigiensis (see AL MABRUK et al., 2021 and GROUD et al., 2021 from Algeria), thus suggesting that it crossed the Suez Canal as a Lessepsian migrant. As a result of the taxonomic complex, it is not clear how many records of this species have been reported from the Mediterranean. GROUD et al., (2021) mentions ten records of Kyphosus spp., however only half of them were identified as K. vaigiensis.

REFERENCES

AL MABRUK, S.A.A., A. ABDULGHANI, O.A.NOUR, M. ADEL, F. CROCETTA, N. DOUMPAS, P. KLEITOU & F. TIRALONGO. 2021. The role of social media in compensating for the lack of field studues: Five new fish species for Mediterranean Egypt. J. Fish Biol., 99(2): 673-678. https://doi.org/ 10.1111/jfb.14721.

AZZURRO, E., L. PENA-RIVAS, D. LLORIS & M.

BARICHE. 2013. First documented occurrence of *Kyphosus incisor* in the Mediterranean Sea. Mar. Biodiver. Rec. https://doi.org/10.1017/S1755267213000717; Vol. 6; e98

FRANCOUR, P. & N. MOUINE. 2008. First record of *Kyphosus spectator* (Linnaeus, 1758) (Kyphosidae) along the French Mediterranean coast. Cybium 32: 275–276.

- GOLANI, D., L.ORSI-RELINI, E. MASSUTI & J.P. QUGIGNARD 2002. *CIESM atlas of exotic species in the Mediterranean*. Vol.1. Fishes. Briand, F. Editor. CIESM Publications, Monaco, 256 pp.
- GOLANI, D., E. AZZURRO, J. DULČIĆ, E. MAS-SUTI & L. ORSI-RELINI. 2021. Atlas of Exotic Fishes in the Mediterranean Sea. 2nd edition [F. Briand, Editor.] 365 pages. CIESM Publishers, Paris, Monaco.
- GOREN, M., B.S. GALIL, R. GEVILI & N. STERN. 2016. First record of the brassy chub *Kyphosus vaigiensis* (Quoy&Gaimard, 1825) in the eastern Mediterranean (Osteichthyes: Perciformes: Kyphosidae). Zool. Midd. East 62: 1–4.
- GROUD, L.L., L. CHAOUI & M.H. Kara. 2021. A new record of the brassy chub *Kyphosus vaigiensis* (Actinopterygii: Perciformes: Kyphosidae) from the Mediterranean Sea. Acta Ichthyolog. Piscat. 51(2): 219-223. https://doi.org/10.3897/aiep.51.64069.
- HUBBS, C.L. & K.F. LAGLER. 1947. Fishes of the Great Lakes Region. Cranbrook Institute of Science. Michigan, USA. 186 pp.
- KIPARISSIS, S., D. LOUKOVITIS & C. BATAR-GIAS. 2012. First record of the Bermuda Sea chub *Kyphosus saltatrix* (Pisces: Kyphosidae) in Greek waters. Mar. Biodiver. Rec. 5: e11.

- KNUDSEN, S.W. & K.D. CLEMENTS. 2013. Revision of the fish family Kyphosidae (Teleostei: Perciformes). Zootaxa 3751(1):1-101
- LIGAS, A., P. SARTOR, M. SBRANA & S. DE RANI-ERI. 2011. A new record of *Kyphosus saltatrix* (Pisces: Kyphosidae) along the Italian coasts (North-Western Mediterranean). Mar. Biodiver. Rec. 4: e6.
- NELSON, J.S., T.C. GRANDE & M.V.H. WILSON. 2016. Fishes of the World. 5th ed. John Wiley & sons. Hoboken, New Jersey, USA. 707 pp.
- ORSI-RELINI, L. 2017. Note on recent revisions of the taxonomy of Kyphosidae. Biol. Mar. Med. 24(1): 206-208.
- ORSI-RELINI, L., M.R. COSTA & M. RELINI. 2010. First record of the yellow sea chub *Kyphosus incisor* in the Mediterranean. Mar. Biodiver. Rec. 3: e4.
- SAKAI, K. 2016. Kyphosidae. Pp. 2680-2685. In: Carpenter, K.E and De Angolis (eds.). The Living Marine Resources of the Eastern Central Atlantic. Vol. 4: Bony fishes part 2 (Perciformes to Tetraodontiformes) and Sea Turtles. FAO Species Identification Guide for Fishery Purposes. Rome, FAO.
- TORTONESE, E. 1986. Kyphosidae. 912-913 pp. In: Whitehead P.J.P., Bauchot M.-L., Hureau J.-C., Nielsen J., Tortonese E. (Editors), Fishes of the North-eastern Atlantic and the Mediterranean. Vol. 2. UNESCO, Paris: 517-1007.

Received:2 November 2021 Accepted:3 May 2022

Prvi zapis o nalazu *Kyphosus vaigiensis* (Ribe: Kyphosidae) na sredozemnoj obale Libije

Amani FITORI, Ali EL-FITURI i Daniel GOLANI

Kontakt e-pošta: dani.golani@mail.huji.ac.il

SAŽETAK

Prvi zapis o nalazu vrste *Kyphosus vaigiensis* (Ribe: Kyphosidae) na sredozemnoj obali Libije. Pretpostavlja se da je ova vrsta u Sredozemlje stigla iz Crvenog mora preko Sueskog kanala. Ističemo potrebu taksonomskog istraživanja roda *Kyphosus* u Sredozemlju.

Ključne riječi: Kyphosus vaigiensis; Sredozemlje; Libija; invazivne vrste