

New data on *Pinctada radiata* (Leach, 1814) (Bivalvia: Pteriidae) in the Adriatic Sea

Slavica PETOVIĆ¹* and Vesna MAČIĆ¹

University of Montenegro, Institute of Marine Biology, Dobrota bb, 85330 Kotor, Montenegro

*Corresponding author: kascelanslavica@gmail.com

The first recorded population of *Pinctada radiata* in the Adriatic Sea was observed on the pier in Porto Montenegro marina (Tivat, Montenegro), in August 2016. Numerous individuals inhabited pier walls at a depth of 5 m. The shell height (SH) values for the 15 randomly collected individuals of *P. radiata* ranged between 32.2 mm and 52.1 mm with an average SH value of 38.3 mm (standard deviation ±6.1). Further surveys in Montenegro will provide information on the establishment of the observed population and what is its impact on local biodiversity.

Key words: alien species, *Pinctada radiata*, morphometric parameters, Adriatic Sea

INTRODUCTION

Pinctada radiata (Leach, 1814) known as ‘pearl oyster’, is widespread in shallow waters of the tropical and subtropical continental shelf regions and particularly abundant in the Indo-Pacific (WADA & TËMKIN, 2008). Shell morphology of this species varies with salinity, usually is up to 50-65 mm but could exceed 100 mm in length and it is protandric hermaphrodite species with sex inversion occurring in shells 32-57 mm (ZENETOS *et al.*, 2003). Furthermore, it is one of three alien species from family Pteriidae spread in the Mediterranean Sea. Other two species are *Pinctada margaritifera* (Linné, 1758) and *Electroma vexillum* (Reeve, 1857) (ÇEVİK *et al.*, 2005; ZENETOS *et al.*, 2010).

Pinctada radiata is considered as the first lessepsian bivalve species reported for the Mediterranean Sea, being described from Egypt in 1884 under the synonym *Meleagrina savignyi* (MONTEROSATO, 1878). Since then the species

has successfully spread throughout the Mediterranean Sea, colonizing new habitats and becoming very abundant in the Levantine Sea, Greece, Libya, Turkey, Syria, Tunisia, Malta and France (DEIDUN *et al.*, 2014). It was also intentionally imported for mariculture in many areas of Greece and Italy during the last century (KATSANEVAKIS *et al.*, 2008). For better understanding on biometric, reproductive, demographic and genetic features of this bivalve species several detailed studies on the Mediterranean populations of *P. radiata* were performed in Tunisia (TLIG-ZOUARI & ZOUALI, 1995; DERBALI *et al.*, 2011; BELLAJJ-ZOUARI *et al.*, 2012), Egypt (YASSIEN *et al.*, 2000), Turkey (GOKOGLU *et al.*, 2006) and Malta (DEIDUN *et al.*, 2014).

In the Adriatic Sea, only few specimens were recorded up to date. VIO & DE MIN (1996) described this species in Trieste Bay as live individuals attached to an oil platform originating from the Sicilian Channel. In the Gulf of Trieste, the species has never been recorded again and

further researches in the area seem to confirm that it did not survive in the area (CROCETTA *et al.*, 2009). Recently, two juvenile specimens (less than 3 mm in shell length) from 59 m on a silty-sand bottom in the Croatian waters were identified as *Pinctada radiata* by DOĞAN & NERLOVIĆ (2008). A very recent record from the Italian Adriatic Sea (Torre Guaceto) was also published (SCUDERI & TERLIZZI, 2012).

Here we present the first record of *P. radiata* population from the Adriatic Sea as a basis for future monitoring of this species, listed as one of the 100 worst invasives (STREFTARIS & ZENETOS, 2006).

MATERIALS AND METHODS

Research was performed in Porto Montenegro marina (Tivat, Montenegro) (Fig. 1) as a part of Biological monitoring program, in August 2016. Fouling material from the piers was surveyed and photographed by SCUBA diving, while some organisms were collected for further analysis in laboratory. In the holes between the concrete blocks of the pier, at 5 m depths, 15 randomly selected individuals of *P. radiata* were collected. Measurements were taken using an electronic digital calliper (precision of 0.01 mm) so that comprised shell height (SH), shell length (SL), hinge length (HL) and shell width (SW) were measured. The material was preserved in 70% ethanol.




Fig. 1. Sampling location of *P. radiata* in Porto Montenegro marina (Tivat, Montenegro)

RESULTS AND DISCUSSION

This record of *P. radiata* is the first one for the south Adriatic and first data about large population in the Adriatic Sea. Values for the shell height (SH) ranged between 32 mm and 52 mm with an average SH value of 38.3 mm (standard deviation ± 6.1) (Table 1). Similar individual size is also found in Maltese island (DEIDUN *et al.*, 2014), while a maximum value reported from other central Mediterranean localities, including Linosa 78.7 mm (LODOLA *et al.*, 2013), El Bibane Lagoon in Tunisia 85.0 mm (SEURAT, 1929), Bizerte Lagoon in Tunisia 100.5 mm (TLIG-ZOUARI *et al.*, 2010), Hammamet in Tunisia 104.3 mm (BELLAAJ-ZOUARI *et al.*, 2012) and the Gulf of Gabes in Tunisia 96.0 mm (DERBALI *et al.*, 2011) is markedly different.

Table 1. Biometric parameters of the *P. radiata* specimens sampled in Porto Montenegro marina, Montenegro

Biometric parameters	Mean values (\pm st dev in mm)
Shell length (SL)	38.3 (± 7.3)
Hinge length (HL)	39.7 (± 5.3)
Shell height (SH)	38.3 (± 6.1)
Shell width (SW)	11.4 (± 2.1)

Studying spatial patterns of marine alien species across the Ionian–Adriatic boundary, KATSANEVAKIS *et al.*, (2011) found one *P. radiata* alive in the north part of Ionian Sea at only a single site (Saranda, Albania). Our record of new population in the south Adriatic Sea, together with the global warming signals (COLL *et al.*, 2010), looks like spreading on the north, as a result of natural dispersal via larval transport by currents, but we should also have in mind possible dispersal by fouling. Interesting is also that *P. radiata* was found in Porto Montenegro marina only on newly constructed parts of the pier 1 (constructed in 2015), confirming its high ability in competition for space with native species (Fig. 2).

Because of its adaptation to subtropical environment and tolerance to chemical contami-




Fig. 2. *Pinctada radiata* in situ (A) and bivalve collected for morphometric analyses (B)

nation (KATSANEVAKIS *et al.*, 2008) it is to expect that this population will survive. Our findings justify the setting up of a monitoring program in Porto Montenegro marina, while further surveys will provide more accurate information on its abundance, distribution in marina (and possibly in the surrounding) as well as possible impact on native biodiversity.

ACKNOWLEDGEMENTS

The study was carried out within the framework of the 2016/2017 biological monitoring program financed by Porto Montenegro Marina, Adriatic Marinas.

REFERENCES

- BELLAJJ-ZOUARI, A., S. DKHILI, R. GHARSALLI, A. DERBALI & N. ALOUI-BEJAOU. 2012. Shell morphology and relative growth variability of the invasive pearl oyster *Pinctada radiata* in coastal Tunisia. J. Mar. Biol. Assoc. U.K., 92(3): 553–563.
- COLL, M., C. PIRODDI, J. STEENBEEK, K. KASCHNER, , LASRAM, FBR, AGUZZI, J., BALLES-TEROS, C., BIANCHI, N., CORBERA, J., DAILIANIS, T. *et al.*,. 2010. The Biodiversity of the Mediterranean Sea: Estimates, Patterns, and Threats. PLoS ONE 5:e11842.
- CROCETTA, F., W. RENDA W. & A. VAZZANA. 2009. Alien Mollusca along the Calabrian shores of the Messina Strait area and a review of their distribution in the Italian Seas. Boll. Malacol 45: 15-30.
- ÇEVİK, C., A. DOĞAN, M. ÖNEN & A. ZENETOS. 2005. First record of the Indo-Pacific species *Electromma vexillum* (Mollusca: Bivalvia: Pterioida) in the eastern Mediterranean. JMBA2 - Biodiversity Records. doi:10.1017/S0025315 4050121 3X
- DEIDUN, A., F. GIANNI, D.P. CILIA, A. LODOLA & D. SAVINI. 2014. Morphometric analyses of a *Pinctada radiata* (Leach, 1814) (Bivalvia: Pteriidae) population in the Maltese Islands. J Black Sea/Mediterranean Environment 20(1): 1-12.
- DERBALI, A., O. JARBOUI, & M. GHORBEL. 2011. Distribution, abundance and population structure of *Pinctada radiata* (Mollusca: Bivalvia) in southern Tunisian waters (Central Mediterranean). Cah Biol Mar 52: 23–31.
- DOĞAN, A. & V. NERLOVIĆ. 2008. On the occurrence of *Pinctada radiata* (Mollusca: Bivalvia: Pteriidae), an alien species in Croatian waters. Acta Adriatica 49 (2): 155-158.
- GOKOGLU, N., M. GOKOGLU. & P. YERLIKAYA. 2006. Seasonal variations in proximate and elemental composition of pearl oyster (*Pinctada radiata*, Leach, 1814). J. Sci. Food Agric. 86: 2161–2165.
- KATSANEVAKIS, S., E. LEFKADITOU, S. GALINOU-MITSOURI, D. KOUTSOURAS & A. ZENETOS. 2008. Molluscan species of minor commercial interest in Hellenic Seas: distribution, exploitation and conservation status. Med Mar Sci 9:77–118.

- KATSANEVAKIS, S., A. ZENETOS, V. MAČIĆ, S. BEQIRAJ. & D. POURSANIDIS. 2011. Invading the Adriatic: spatial patterns of marine alien species across the Ionian–Adriatic boundary. *Aquat. Biol.*, 13:107–118.
- LODOLA, A., L. NICOLINI, D. SAVINI, A. DEIDUN. & A. OCCHIPINTI-AMBROGI. 2013. Range expansion and biometric features of *Pinctada imbricata radiata* (Bivalvia: Pteriidae) around Linosa Island, Central Mediterranean Sea (Italy). *Ital J Zool* 80(2), 303-312.
- MONTEROSATO, T.A. 1878. Enumerazione e sinonimia delle conchiglie mediterranee. *Gior-*
nale di Scienze Naturali ed Economiche di Palermo 13:61-115.
- SEURAT, L.G. 1929. Observations sur les limites, les faciès et les associations animales de l'étage intercotidal de la petite Syrte (Golfe de Gabès). *B. Inst. Nat. Sci. Tec. Mer. Salammbô* 3:1–71.
- SCUDERI, D. & A. TERLIZZI 2012. Manuale di Malacologia dell'Alto Jonio. Lecce: Edizioni Grifo, 188 pp. in STASOLA, G., RIOLO, F., MACALI, A., PIERRI, C. & F. CROCETTA. 2014. Further spreading in the Italian seas of already established non-indigenous mollusk species. *Marine Biodiversity Records*, Vol. 7: 1-8.
- STREFTARIS, N. & A. ZENETOS. 2006. Alien marine species in the Mediterranean- the 100 “Worst Invasives” and their impact. *Med Mar Sci* 7(1):87-118.
- TLIG-ZOUARI, S., L. RABAQUI, I. IRATHNI, M. DIAWARA & O. K. BEN HASSINE. 2010. Comparative morphometric study of the invasive pearl oyster *Pinctada radiata* along the Tunisian coastline. *Biol* 65(2): 294-300.
- TLIG-ZOUARI, S. & J. ZAOUALI. 1995. La population de *Pinctada radiata* des îles Kerkennah (Tunisie). Paramètres démographiques et croissance. *B Soc Sci Nat Tun* 24: 103-117.
- VIO, E. & R. DE MIN. 1996. Contributo alla conoscenza dei Molluschi marini del Golfo di Trieste. *Atti Mus Civ Stor Nat Trieste* 47: 173-233.
- WADA, K.T. & I. TËMKIN. 2008. Taxonomy and phylogeny. In: Southgate P C, Lucas J S. (Editors). Elsevier, Philadelphia, USA., p. 37-75.
- YASSIEN, M.H., F. A. ABD EL-RAZEK & R. KILADA. 2000. Growth estimates of the Pearl Oyster, *Pinctada radiata*, from the eastern Mediterranean. *Egypt J Aq Biol Fish* 4:105–118.
- ZENETOS, A., S. GOFAS, G., RUSSO & J. TEMPLADO. 2003. CIESM Atlas of Exotic Species in the Mediterranean. Vol. 3. Molluscs. In: F. Briand (Editor). CIESM Publishers, Monaco, 376 pp.
- ZENETOS, A., S. GOFAS, M. VERLAQUE, M.E. INAR, J.E. GARCIA RASO, C.N. BIANCHI, C. MORRI, E. AZZURRO, M. BILECENOGLU, C. FROGLIA. 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union’s Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Med Mar Sci* 11(2): 381–493.

Received: 12 December 2016

Accepted: 26 July 2017

Novi podaci o vrsti *Pinctada radiata* (Leach, 1814) (Bivalvia: Pteriidae) u Jadranskom moru

Slavica PETOVIĆ* i Vesna MAČIĆ

*Kontakt e-adresa: kascelanslavica@gmail.com

SAŽETAK

Prvi nalazi populacije *Pinctada radiata* u Jadranskom moru se odnose na dokove marine Porto Montenegro (Tivat, Crna Gora), u kolovozu 2016. godine. Na zidovima dokova zabilježene su brojne jedinke na dubini od 5m. Visina ljuštture (SH) za 15 slučajno odabranih jedinki *P. radiata* je kolebala od 32.2 mm do 52.1 mm sa srednjom vrijednošću od 38.3 mm (stand. dev. ± 6.1). Daljna istraživanja u Crnoj Gori će omogućiti informacije o uspješnosti uspostavljanja ove populacije i njenom uticaju na lokalnu bioraznolikost.

Ključne riječi: unsene vrste, *Pinctada radiata*, morfometrijski parametri, Jadransko more

