

Occurrence of two rare species from order *Lampriformes*: Crestfish *Lophotus lacepede* (Giorna, 1809) and scalloped ribbonfish *Zu cristatus* (Bonelli, 1819) in the northern coast of Sicily, Italy

Fabio FALSONE¹, Michele Luca GERACI¹, Danilo SCANNELLA¹,
Charles Odilichukwu R. OKPALA¹, Giovan Battista GIUSTO¹, Mar BOSCH-BELMAR²,
Salvatore GANCITANO¹ and Gioacchino BONO¹

¹*Institute for the Coastal Marine Environment, IAMC-CNR,
91026 Mazara del Vallo, Sicily, Italy*

²*Consorzio Nazionale Interuniversitario per le Scienze del Mare (CoNISMa), Rome, Italy*

Corresponding author, e-mail: falsonefabio@gmail.com

The bony fish Lophotus lacepede (Giorna, 1809) and Zu cristatus (Bonelli, 1819) are the two species rarely recorded within the Mediterranean basin, usually reported as accidentally captured in depth (mesopelagic) fishing operations. In the current work, we present the first record of L. lacepede and Z. cristatus in fishing catches from southwestern Tyrrhenian Sea. Moreover, in order to improve existent biological/ecological knowledge, some bio-related aspects such as feeding aspect, sexual maturity and age estimate have been discussed.

Key words: crestfish, scalloped ribbonfish, meristic features, vertebrae, growth ring

INTRODUCTION

The target species of this study (*Lophotus lacepede* and *Zu cristatus*) belong to Lophotidae (Bonaparte, 1845) and Trachipteridae (Swainson, 1839) families respectively, including the Lampriformes order (consisted of 7 families).

L. lacepede (largely referred to as crestfish) is a pan-oceanic species, which can reach up to 200 cm as maximum total length (TL). Despite its wide distribution, it is rare to find and its catches appear to be widely separated from each other by either space or time (GASTON, 1994). Being the only Mediterranean representative

species of Lophotidae family, the *L. lacepede* inhabits the epipelagic zone, although it could also be recorded in most oceans from the surface up to 1000 m depth (HEEMSTRA, 1986; PALMER, 1986; OLNEY, 1999). First record of this species in the Mediterranean Basin was from the Southern Adriatic Sea (KOLOMBATOVIĆ, 1890) and afterwards it was observed in other places, ranging from the Aegean Sea to the Spanish coast (MINOS *et al.*, 2015), as well as, off the coast of Algeria (BACHOUCHE *et al.*, 2016). Essentially, this species derives its name from the occipital adipose crest situated on top of the head holding up vertically to the first ray of dorsal fins




Fig. 1. Map of southwestern Tyrrhenian Sea location (full orange triangle) sampling of *L. lacepede* and *Z. cristatus*

(JARDAS, 1996). Besides, juveniles of this species feed on plankton while adults feed on either fish and/or squids (REY, 1983; MINOS *et al.*, 2015).

On the other hand, Trachipteridae family includes three genera *Desmodema* (Walters & Fitch, 1960) *Trachipterus* (Goüan, 1770) and *Zu* (Walters & Fitch, 1960). *Z. cristatus* (also called scalloped ribbonfish), well-known due to its particular face-up swimming (BIANCHI *et al.*, 1999), seems to be the only Mediterranean species from its genus. Furthermore, it is a cosmopolitan species, since it occupies ocean waters from tropical to temperate latitudes (FROESE & PAULY, 2005). Regarding the bathymetric distribution, it might be considered as mesopelagic, given its existence in wider depths of up to 800 m. Mostly, scalloped ribbonfish adults tend to feed on small fish and cephalopods (PALMER, 1986). In the Mediterranean basin, records of *Z. cristatus* are relatively rare (BIANCO *et al.*, 2006) and its capture has long been seen to mostly concur with the principal spawning period within its seasonal life cycle (SPARTA, 1956). In particular, the occurrence of *Z. cristatus* has been reported in the Adriatic and Ligurian Sea, central and south-eastern Tyrrhenian Sea, off the coasts of Spain and Algeria and in the Gulf of Tunis (north of Tunisia) (FISCHER, *et al.*, 1987; PAPACOSTANTINOU, 1988; QUIGNARD & TOMASINI, 2000; GOLANI *et al.*, 2006; ZENETOS *et al.*, 2015).

In the current work, an unexpected catch of *L. lacepede* and *Z. cristatus* species at south-

western Tyrrhenian Sea has been described, together with some aspects of the species biology such as feeding, sexual maturity and age estimation.

MATERIAL AND METHODS

During the summer 2015 season, two fish specimens from order Lampriformes, (*L. lacepede* and *Z. cristatus*) were caught using a long line drifting 20 miles off San Vito Lo Capo, Sicily (38°29'00"N, 12°31'24"E). Afterwards, samples were transported to the IAMC-CNR laboratory in Mazara del Vallo where individuals were examined and biometric and meristic data were collected. Biometrics included various lengths, widths and weights of different fish body parts. Meristic data was referred to numerical counts of various fins, vertebrae, teeth and gill rakers. All measurements were performed by visual inspection using available expertise. Collected data was expressed in terms of millimeter (mm) and subsequently as percentage (%) of standard length (SL) while all weights were measured with an accuracy of 0.1 g.

The use of photographs was employed to supplement all observed data. This was performed prior to and after dissections, which were intended to establish features such as age estimations, sex of fish, contents of the stomach, as well as internal organs. For instance, to estimate the age, two independent experts identified and counted the accretion bands (annual growth rings that allow for age estimation) by means of hard structures. To further support age estimation, the second vertebra has been used at both species adding the first dorsal ray (*L. lacepede*) and otolith (*Z. cristatus*). To define the sexual maturity (SM), the MEDITS (International Bottom Trawl Survey in the Mediterranean) approach has been employed by way of visual inspection of gonads. In particular, sexual maturity has been classified as follows: Stage 0 - undetermined; Stage 1 - immature; Stage 2a - virgin-developing; Stage 2b - recovering; Stage 2c - maturing; Stage 3 - mature/spawner; Stage 4a - spent; as well as Stage 4b - resting (BERTRAND *et al.*, 2007). All measurements were performed using preserved specimens.

In order to perform robust comparisons of our current data, we used comparable data records of these species already reported in other Mediterranean areas such as the Strait of Sicily (RAGONESE *et al.*, 1997), Adriatic Sea (ŠPREM *et al.*, 2014; DULČIĆ, 2002) and marine waters of Algeria (BACHOUCHE *et al.*, 2016) for *L. lacepede*, while the Ligurian Sea (TORTONESE, 1958; PSOMADAKIS *et al.*, 2007) and the Iberian Sea (ROIG DEMESTRE, 1982) for *Z. cristatus*.

RESULTS AND DISCUSSION

The biometric data obtained during the study were similar to those previously reported by other authors (Table 1) (BACHOUCHE *et al.*, 2016; TORTONESE, 1958; RAGONESE *et al.*, 1997; ŠPREM *et al.*, 2014; PSOMADAKIS *et al.*, 2007; ROIG & DEMESTRE, 1982).

Regarding *L. lacepede*, only the operculum height resulted higher compared to the data reported by RAGONESE *et al.* (1997), 74.8% and 62.8% of the SL respectively. Concerning *Z. cristatus*, just the dorsal fin maximum height was higher in comparison to the data recorded by PSOMADAKIS *et al.* (2007), 12.5 % and 6.6% of the SL respectively. To our best knowledge, in the Mediterranean Sea, information regarding the operculum thickness, gonad length, as well as weights of stomach content/empty, intestine, and gonad, ink gland/duct of *L. lacepede* was not available in the literature. Similarly, specific data on weights of stomach content/empty, liver and gonad, as well as other measurements such as operculum thickness/heights, dorsal fin (maximum length), base pectoral fin (width), gonad length, number of vertebra, and teeth (lower/




Fig. 2. Images of the whole, head, gonad, one vertebra specimen and transverse section of first fin dorsalray for *L. lacepede*

b) MERISTIC (COUNTS)	<i>Lophotus lacepede</i>			<i>Zu cristatus</i>			
	Present note	Strait of Sicily ¹	Algerian coast ³	Present note	Ligurian sea ⁴	Ligurian sea ⁵	Iberian sea ⁶
Number rays dorsal fin	236	248	228	126	130	125	117
Number rays pectoral fin	17	16	15	11	11	10	11
Number rays anal fin	17	20	20				
Number rays caudal fin		18	19	9	9	9+1	9+4
Number vertebrae	136	140		62			
Gill rakers ***				3+8	3+8	2+9	10
Teeth upper jaw				18	21		
Teeth lower jaw				12	10		
Palatin teeth				4	4		
Vomerin teeth				3	4		

¹ Literature from RAGONESE *et al.*, 1997; ² Literature from ŠPREM *et al.*, 2014; ³ Literature from BACHOUCHE *et al.*, 2016; ⁴ Literature from PSOMADAKIS *et al.*, 2007; ⁵ Literature from TORTONESE 1958; ⁶ Literature from ROIG & DEMESTRE 1982.

% SL was referred as a percentage of the standard length; % TL was referred as a percentage of the total length; % TW was referred as a percentage of the total weight

*** The first number was referred to the epibranchial and the second number to ceratobranchial (first left gill arch)

upper jaw/palate/vomer) of *Z. cristatus* have not been earlier reported.

In addition, a thorough visual inspection of samples' anatomo-morphological features was conducted. Images of the whole, head, gonad, one vertebra specimen and a transverse section of first fin dorsal ray for *L. lacepede* are depicted in Fig. 2. Also, different pictures of the whole, head, gonad, one vertebra specimen and transverse section of otolith for *Z. cristatus* are included in Fig. 3.

The diagnostic identification, based on visual observations, showed that *L. lacepede* sample was a female whereas *Z. cristatus* was a male. Thereafter, both specimens were dissected to allow for further analyses. For *L. lacepede*, the gonad length and weight were \approx 613 mm and 42 g, respectively. The ovaries seemed "Y" shaped and were well developed at maturity stage 3 (gonads occupied more than two-thirds of the abdominal cavity), indicating that it was a mature female. The calculated Gonadosomatic Index (GSI), which represents gonad as the proportion to fish body mass widely believed to express the fish maturity associative of development of ovary/testes, has been estimated at

1.77. For *Z. cristatus*, the testicles length and weight were \approx 146 mm and 3 g respectively, which seemed well developed at maturity stage 3. However, the GSI was 0.24 that it is a low value if compared to the recognized maturity stage. This information on gonads (ovaries/testes) and the period of capture, may offer some insights about the reproduction of inspected samples. In the present study, in agreement with previous literature, the spawning period of *L. lacepede* seemed related to the summer season (MINOS *et al.*, 2015; SANZO 1940; SPARTÀ 1954). However, DULČIĆ & SOLDI (2008), found a mature spawner male during the winter period. Regarding *Z. cristatus*, previous studies reported that the spawning period of this species usually takes place between spring and summer seasons (DULČIĆ, 2002; PSOMADAKIS *et al.*, 2007, BRADAI & OUAER, 2012); nevertheless, *Z. cristatus* eggs have been collected during summer–fall period at Middle Atlantic Bight (OLNEY, J.E. & A. NAPLIN 1980). Notably, DULČIĆ (2002) reported that the preferred spawning ground of *Z. cristatus* was related to low salinity environments. However, the current study may suggest that these species would tolerate wider salinity ranges since



Fig. 3. Images of the whole, head, gonad, one vertebra specimen and transverse section of otolith for *Z. cristatus*

examined specimens were collected at open sea (38%). At the light of this, additional information would be necessary to better understand the factors influencing the spawning period of these species.

To estimate the age of *L. lacepede*, seven accretion rings were counted from both vertebrae and dorsal ray sections, while for *Z. cristatus*, eleven accretion rings were counted from otolith transverse section and vertebrae. Whilst

accretion rings fundamentally accord with the age of fish (PANFILI *et al.*, 2002; PRINCE & PULOS, 1983) and although observations herein have been performed based on one specimen, this age estimation together with the maturity stage further indicates that the studied samples corresponded with adult individuals. Heretofore, this should represent the first attempt to estimate the age of these species within the Mediterranean basin.

The total stomach contents of *L. lacepede* were also analyzed. We found (a) five specimens of *Scaevargus unicirrhus* (Delle Chiaje [in Férussac & d'Orbigny], 1841) (b) one specimen of *Sepietta oweniana* d'Orbigny, 1841 (c) thirteen cephalopods since twenty-five mandibles were counted, (d) remains of Portunidae paws and (e) remains of skeleton and scales fish. It thereafter allowed for a classification, namely: 93.3% of Cephalopoda class (a, b, c respectively), 1.4% of Malacostraca class (d) and 5.3% of Osteichthyes class (e). Possibly, *L. lacepede* diet has been mainly based on cephalopods followed by bony fish and crustaceans somewhat in agreement with MINOS *et al.* (2015). The stomach of *Z. cristatus* contained only the head of Scombridae fish, which probably was used as the bait of long-line fishery.

CONCLUSIONS

To the best of our knowledge, this is the first report about *L. lacepede* and *Z. cristatus* species unexpectedly caught off the southwestern Tyrrhenian Sea. Previous authors such as TORTONESE (1958) described these species as rarely captured probably because of reduced fishing activity that took place at mesopelagic habitats at those times. The present study contributes to improving the extant biological knowledge, such as the feeding aspect, sexual maturity and age estimation of these species in the Mediterranean basin. Finally, we would like to point out that further investigation should be conducted to better understand the biology and life cycle of these rare species.

REFERENCES

- BACHOUCHE S., M. ETSOURI, S. ROUIDI, B. COLONEL & A.B. TIPAZA. 2016. The first record of Crested Oarfish, *Lophotus lacepede* (Actinopterygii: Lampriformes) from the marine waters of Algeria. *Int. J. Sci. know.*, 1: 19-22.
- BERTRAND J., SOUPLET A., GIL DE SOLA L., RELINI G. & CY POLITOU. 2007. International bottom trawl survey in the Mediterranean (MEDITS). *MEDIT Survey, Instruction manual, version 5*. IFREMER. Nantes cedex France, 62 pp.
- BIANCHI, G., K.E. CARPENTER, J.P. ROUX, F.J. MOLLOY, D. BOYER & H.J. BOYER. 1999. *FAO species identification guide for fishery purposes. Field guide to the living marine resources of Namibia*. FAO, Rome, 265 pp.
- BIANCO, P.G., V. ZUPO & V. KETMAIER. 2006. Occurrence of the scalloped ribbonfish *Zu cristatus* (Lampridiformes) in coastal waters of the central Tyrrhenian Sea, Italy. *J. Fish Biol.*, 68:150-155.
- BRADAI, M.N. & A. EL OUAER. 2012. New record of the scalloped ribbon fish, *Zu cristatus* (Osteichthyes: Trachipteridae) in Tunisian waters (central Mediterranean). *Mar. Biodivers. Rec.*, 5: 1-3.
- DULČIĆ, J. & H. AHNELT. 2007. How many specimens of the crested oarfish, *Lophotus lacepede* Giorna, 1809 (Pisces: Lophotidae), were caught in the Adriatic Sea? *Acta Adriat.*, 48: 39-43.
- DULČIĆ, J. 2002. First record of scalloped ribbon fish, *Zu cristatus* (Pisces: Trachipteridae), eggs in the Adriatic Sea. *J. Plankton Res.*, 24: 1245-1246.
- FISHER, W., M.L. BAUCHOT. & M.S. SCHNEIDER. 1987. *Fiches FAO d'Identification des Espèces pour les Besoins de la Pêche. Méditerranée et Mer Noire, vol. 2 (Vertébrés)*. FAO, Rome, 2:761-1529.
- GASTON, K.J. 1994. What is rarity?. In: Chapman and Hall (Editors). *Population and Community Biology. Series, vol. 13*, New York, N.Y., 205 pp.
- GOLANI, D., B. ÖZTÜRK, N. BAŞUSTA. 2006. *Fishes of the Eastern Mediterranean*. Turkish Marine Research Foundation, Publication no.24, Istanbul, Turkey, 259 pp.
- HEEMSTRA, P.C. 1986. Lophotidae. In: M.M. Smith and P.C. Heemstra (Editors). *Smiths' sea fishes*, Springer-Verlag: Berlin, 402-403.
- JARDAS, I. 1996. The Adriatic ichthyofauna (in Croatian with English summary). *Školska knjiga d. d., Zagreb*, 533 pp.
- KOLOMBATOVIĆ, J. 1890. *Notizie ittologiche*. In:

- Glas. Hrvat. Naravosl. Druš., 5:167-168.
- MINOS, G., S.T. KARIDAS & P.S. ECONOMIDIS. 2015. Range extension for crested oarfish *Lophotus lacepede* Giorna, 1809 in the waters of the northern Aegean Sea, Greece. *Acta Adriat.*, 56(2): 283-290.
- OLNEY, J. E. & A. NAPLIN. 1980. Eggs of the Scalloped Ribbonfish, *Zu cristatus*, (Pisces: Trachipteridae) in the Western North Atlantic. *Copeia*, 1: 165-166.
- OLNEY, J.E. 1999. Lophotidae. The living marine resources of the Western Central Pacific. In: K.E. Carpenter & V.H. Niem (Editors), vol. 3: Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae). FAO, Rome: 1970-1971.
- PALMER, G. 1986. Lophotidae. In: P.J.P. Whitehead, M.L. Bauchot, J.C. Hureau, J. Nielsen and E. Tortonese (Editors). *Fishes of the north-eastern Atlantic and the Mediterranean*. Volume 2. Richard Clay (The Chaucer Press) Ltd, Bungay (United Kingdom): 734-735.
- PANFILI, J., H. DE PONTUAL, H. TROADEC & P.J. WRIGHT. 2002. *Manual of fish sclerochronology*. Brest, France: Ifremer-IRD coedition, 466 pp.
- PAPACONSTANTINO, C. 1988. *Fauna Graeciae, IV. Check-list of marine fishes of Greece*. NCMR, Hellenic Zoological Society, 257 pp.
- PRINCE E.D. & L.M. PULOS. 1983. Preface. *Proceedings of the International Workshop on Age Determination of Oceanic Pelagic Fishes: Tunas, Billfishes and Sharks*. US Dep. Commer. NOAA Tech. Rep. NMFS-8.
- PSOMADAKIS P.N., M. BOTTARO, M. VACCHI. 2007. On two large specimens of *Zu cristatus* (Trachipteridae) from the Gulf of Genoa (NW Mediterranean). *Cybium*, 31: 480-482.
- QUIGNARD J. P. & J. A. TOMASINI. 2000. Mediterranean fish biodiversity. *Biol. Mar. Mediterr.* 7: 1-66.
- RAGONESE, S., P. JEREB & U. MORARA. 1997. Su di un esemplare di pesce liocorno *Lophotus lacepedei* (Lampridiformes-Lophotidae) spiaggiato a Mazzara del Vallo (Sicilia Sud Occidentale). (About one specimen of crested oarfish *Lophotus lacepedei* (Lampridiformes-Lophotidae) found at Mazzara del Vallo (southwestern Sicily)). *Biol. Mar. Mediterr.*, 4: 551-553.
- REY, J.C. 1983. Captura de un ejemplar de *Lophotus lacepedei* Giorna, 1809 (Pisces, Lophotidae) en el estrecho de Gibraltar. *Memórias do Museu do Mar - Série Zoológica*, vol. 3(25): 1-16.
- ROIG A. & M. DEMESTRE. 1982. Sobre la captura de dos *Zu cristatus* (Bonelli, 1820) en aguas del litoral catalán (*Pisces, Trachipteridae*). (About catch of two *Zu cristatus* (Bonelli, 1820) in catalan waters (*Pisces, Trachipteridae*)). *Misc. Zool.*, Barcelona, 6: 152-154.
- SPARTA A. 1956. Trachipteridae. *Fauna e flora del Golfo di Napoli*. Monografia. Stazione Zoologica di Napoli, 38: 266-275.
- ŠPREM, J.S., T. DOBROSLAVIĆ, V. KOŽUL, I. PRUSINA, V. ONOFRI & N. ANTOLOVIĆ. 2014. New record of *Lophotus lacepede* (Giorna, 1809) and *Lampris guttatus* (Brünnich, 1788) in the southeastern Adriatic Sea (Croatian coast). *Cah. Biol. Mar.*, 55: 371-373.
- TORTONESE E. 1958. Cattura di *Trachipterus cristatus* (Bonelli). Note sui Trachipteridae del Mar Ligure. *Doriana*. 2:1-5.
- ZENETOS A., AKEL E.H.KH., APOSTOLIDIS C., BILECENOGLU M., BITAR G, BUCHET V., CHALARI N., CORSINI-FOKA M., CROSETTA F., DOGRAMMATZI A., DRAKULIĆ M., FANELLI G., GIGLIO G., IMSIRIDOU A., KAPIRIS K., PKARACHLE P.K., KAVADAS S., KONDYLATOS G., LEFKADITOU E., LIPEJ L., MAVRIČ B., MINOS G., MOUSSA R., PANCUCCI-PAPADOPOULOU M.A., PRATO E., RENDA W., RÍOS N., RIZKALLA S.I., RUSSO F., SERVONNAT M., SIAPATIS A., SPERONE E., THEODOROU J.A., TIRALONGO F., TZOVENIS I. 2015. New Mediterranean Biodiversity Records (April 2015). *Med. Mar. Sci.*, 16 (1): 266-284.

Pojava dviju rijetkih vrsta iz reda *Lampriformes*: britke jedroglavke *Lophotus lacepede* (Giorna, 1809) i vlasuljke *Zu cristatus* (Bonelli, 1819) kod sjeverne obale Sicilije, Italija

Fabio FALSONE, Michele Luca GERACI, Danilo SCANNELLA,
Charles Odilichukwu R. OKPALA, Giovan Battista GIUSTO, Mar BOSCH-BELMAR,
Salvatore GANCITANO i Gioacchino BONO

Kontakt e-adresa: falsonefabio@gmail.com

SAŽETAK

Koštunjičave ribe *Lophotus lacepede* (Giorna, 1809) i *Zu cristatus* (Bonelli, 1819) dvije su vrste koje se rijetko nalaze u Sredozemnom moru i najčešće su zabilježene kao slučajni ulov pri ribolovu dubinskim alatima (mezopelagijal). U ovom radu predstavljamo prvi zapis britke jedroglavke i vlasuljke koje su uhvaćene tijekom ribolova u jugozapadnom dijelu Tirenskog mora. Štoviše, kako bi se poboljšalo postojeće znanje o biologiji i ekologiji, u radu se raspravlja o nekim biološkim određenim aspektima kao što su prehrana, spolna zrelost i procjena starosti.

Ključne riječi: jedroglavka, vlasuljka, meristička obilježja, kralješci, naraštajni prstenovi

