

Report on Cymothoids (Crustacea, Isopoda) collected from marine fishes in Turkey

Ahmet ÖKTENER¹ and Jean-Paul TRILLES²

¹*Cihannüma mah. Hüsnü savman sokak no: 22/5, 80690 Beşiktaş İstanbul, Turkey*

²*UMR 5171 (Génome, Populations, Interactions, Adaptation), Équipe Adaptation Écophysiologique et Ontogenèse, Université de Montpellier 2, CC. 092, Place E. Bataillon, 34095 Montpellier Cedex 05, France*

This work describes Cymothoidae species in Turkey. Some species were collected by the authors, i.e., Anilocra frontalis Milne Edwards, 1840; Nerocila orbignyi (Guérin-Méneville, 1829-1832); Ceratothoa italica Schioedte and Meinert, 1883; Ceratothoa oestroides (Risso, 1826); Ceratothoa parallela (Otto, 1828); Emetha audouini Milne Edwards, 1840; Livoneca punctata (Uljanin, 1872); Mothocya epimerica Costa, 1851, and Mothocya belonae Bruce, 1986. Four of these are reported amongst the Turkish fauna for the first time, i.e., A. frontalis, C. italica, L. punctata, and M. belonae. A list of Cymothoidae already reported from marine fishes of Turkey plus our new reports includes thirteen species (two yet undetermined) from six genera Anilocra, Nerocila, Ceratothoa, Emetha, Livoneca, and Mothocya, and 27 host species. The distributions, hosts, and reporting authors are given.

Key words: Cymothoidae, Crustacea, parasitic isopods, Turkey

INTRODUCTION

Cymothoids (Crustacea, Isopoda) are ectoparasites of marine, freshwater, or brackish water teleost fishes. Numerous families and species of fishes, including many of commercial importance, are infected by cymothoids. They settle on the skin, in the buccal cavity, in the gill chambers, and sometimes in a pouch (TRILLES, 1969; BRUSCA, 1981). Cymothoids are protandrous hermaphrodite isopods with a short phase as free-living, plankton organisms (TRILLES, 1969; COLORNI *et al.*, 1997). Upon finding a suitable

host, they initiate their parasitic life, feeding on blood and tissues (TRILLES, 1969; ROMESTAND & TRILLES, 1976, 1977; ROMESTAND, 1978, 1979; HORTON & OKAMURA, 2003). Several cymothoid isopods have been reported parasitizing marine fishes from the coasts of Turkey (MONOD, 1931; DEMIR, 1952; GELDIAY & KOCATAS, 1972; TRILLES, 1977; AKMIRZA, 1997, 1998a,b, 2000a,b, 2001; KIRKIM, 1998; YELEK, 1998; TOKSEN, 1999; ÖKTENER & SEZGIN, 2000; HORTON & OKAMURA, 2001 & 2003). This study provides a comprehensive summary of Cymothoidae species found in Turkish fauna.

MATERIALS AND METHODS

Three hundred eighty-eight samples belonging to several fish families and species (*Syphodus tinca*, *Spicara maena*, *Dicentrarchus labrax*, *Boops boops*, *Sardina pilchardus*, *Belone belone*, *Alosa fallax*, *Liza aurata*, *Mullus surmuletus*, *Atherina boyeri*) from the Black Sea, the Sea of Marmara, and the Aegean coast of Turkey were examined during 2000-2003. Fish samples were captured with a fish net. The specimens were preserved in 70% alcohol. Data on parasites and collection were recorded. The taxonomic positions of the parasites were identified mainly according to BORCEA (1933), TRILLES (1965, 1968, 1972, 1976, 1977, 1979), TRILLES *et al.* (1989), and BRUCE (1986). The overall prevalence and location of the parasites on their hosts were specified and information about their geographical distribution and host species were added. All collected specimens were deposited in the personal collection of A. ÖKTENER (Cihannüma mah. Hüsnü savman sokak no: 22/5, 80690 Beşiktaş İstanbul, Turkey).

After describing the species that we personally collected, including four species reported for the first time in Turkish fauna, we compiled a list of Cymothoidae already reported for marine fishes of all Turkish coasts. The compilation was prepared as in MOREIRA & SADOWSKY (1978) for the parasitic Isopoda of Chondrichthyes, SIRIKANCHANA (1982) for parasites of fishes in Thailand, and HOLLAND & KENNEDY (1997) for helminth and crustacean species from freshwater fish in Ireland. The following plan was followed: (1) a list of collected Cymothoids and their parasitized fish, the collection site, and author; (2) a list of parasitized fishes with the corresponding species of Cymothoidae.

RESULTS

Cymothoidae collected by the authors

Anilocra Leach, 1818; *Anilocra frontalis* Milne Edwards, 1840

Material:

Anilocra frontalis was collected from the nasal region of two *Syphodus tinca* among

twenty specimens examined (prevalence = 10%) at Sinop (on the Black Sea coast of Turkey). The species is recorded for the first time in the Turkish fauna.

Distribution:

North Sea, Channel, Atlantic Ocean, Mediterranean Sea, Adriatic Sea (TRILLES, 1994).

Hosts:

Especially Labridae (*Labrus maculatus*, *L. vetula*, *L. bergylta*, *L. merula*, *Crenilabrus melops*, *C. cinereus*, *C. ocellatus*); sometimes collected on other fishes (*Gadus* sp., *Merlangius pollachius*, *Blennius pholis*, *Cottus bubalis*, *Gobius flavescens*, *G. paganellus*, *G. minutus*, *Onos mustela*, *Spinachia vulgaris*, *Boops boops*, *B. salpa* and *Spondyliosoma cantharus* (TRILLES, 1994).

Nerocila Leach, 1818; *Nerocila orbignyi* (Guérin-Meneville, 1829-1832)

Material:

Nerocila orbignyi was collected from the operculum on two of four *Liza aurata* (prevalence = 50%) at Sinop.

Distribution:

Australian coasts from central New South Wales, southeastern Tasmania, Victoria, south Australia, and western Australia to Fremantle. Apparently absent from the tropical Indo-Pacific (except for one specimen from Eilat), north Pacific, east Pacific and western north Atlantic (BRUCE, 1987). Widely distributed in the Mediterranean, northwest Africa, Red Sea, ?Egypt, ?New Zealand (TRILLES, 1994).

Hosts:

TRILLES (1975, 1994) summarized the known hosts of this species which chiefly parasitize Mugilidae (*Mugil auratus*, *M. cephalus*, *M. capito*, *M. cheloi*, *M. labrosus*); are less common on *Flesus passer*, *Dicentrarchus labrax*, *Alosa fallax nilotica*, Siphonostomids and Serranidae, ?*Tilapia galilea* and ?*Salmo trutta*. In Australia, BRUCE (1987) recorded the parasite

on *Callorhynchus mihi*, *Acanthopagrus australis*, *Chrysophrys auratus*, *Pseudocaranx dentex*, *Mola mola*, *Sillago bassensis*, *Pomatomus saltatrix* and unidentified Mugilidae, Kyphosidae, and Platyccephalidae, *Girella tricuspidata*, *Chelidonichthys kumu*, and *Dactyloptena orientalis*. *N. orbignyi* is apparently widely ubiquitous but, maybe, several data concern other species.

Ceratothoa Dana, 1852; Ceratothoa italicica
Schioedte et Meinert, 1883

Material:

Ceratothoa italicica was collected from the mouth on seven of thirty-four specimens of *Dicentrarchus labrax* (prevalence = 20.6%) in a sea bass farm on the Mugla coast of the Aegean Sea, Turkey, and from the mouth on four of thirty-five *Spicara maena* (prevalence = 11.4%) in the Bosphorus (Sea of Marmara). *C. italicica* was recorded for the first time in Turkish fauna.

Distribution:

Mediterranean Sea, Adriatic Sea, northwest Africa (TRILLES, 1994; HORTON, 2000).

Hosts:

Pagellus mormyrus, *P. erythrinus*, *Oblada melanura*, *Cantharus lineatus*, *Sargus* sp. (TRILLES, 1994; HORTON, 2000). *D. labrax* and *S. maena* are new hosts, identified for the first time for *C. italicica*.

***Ceratothoa oestroides* (Risso, 1826)**

Material:

Ceratothoa oestroides was collected from the mouth of twelve of forty-five *Spicara maena* (prevalence = 26.7%) in the Erdek coast and the Bosphorus (Sea of Marmara). It was also collected on one of twenty-two *Sardina pilchardus* (prevalence = 4.5%) along the Erdek coast.

Distribution:

Widely distributed in the Mediterranean and Adriatic Seas. Also recorded from the northeast Atlantic including the northwest coasts of Africa (TRILLES, 1994; HORTON, 2000).

Hosts:

This species is found on many host species. Centracanthidae and Sparidae are the most frequently parasitized groups (*Spicara* sp., *S. chrysopis*, *S. maena*, *Smaris melanurus*, *Boops boops*, *Diplodus annularis*, *D. sargus*, *D. vulgaris*). Also, but rarely, collected on *Sardina pilchardus*, *Trachurus trachurus*, *Phycis mediterranea*, *Mullus barbatus*, *Abudebdouf saxatilis* (TRILLES, 1994). Recently recorded from *Dicentrarchus labrax* and *Sparus auratus* on farms of the Adriatic and Greece (PAPOUTSOGLOU et al., 1996; ŠARUŠIĆ, 1999; HORTON & OKAMURA, 2001; MLADINEO, 2002).

***Ceratothoa parallela* (Otto, 1828)**

Material:

Ceratothoa parallela was collected from the mouth of thirteen of twenty-eight *Boops boops* (prevalence = 46.4%) along the Erdek coast (Sea of Marmara).

Distribution:

Widely distributed in the Mediterranean and Adriatic Seas. Also recorded in the northeast Atlantic including the northwest coasts of Africa, the Canary and Madeira Islands (TRILLES, 1994).

Hosts:

Ceratothoa parallela has been found parasitizing particularly *Boops boops*. It was rarely collected on *Sparus*, *Dentex vulgaris*, *Boops salpa*, *Esocis belonis*, *Spicara chrysopis*, *S. maurii*, *S. alcedo*, *Trigla corax*, *Mullus*, *Gadus capelanus*, *Merluccius merluccius*, *Raja asterias*, and *R. clavata* (TRILLES, 1994). However, it seems likely that several reports misidentified the species. Recently, *C. parallela* was reported on cultured *Sparus aurata* in Greece (PAPAPANAGIOTOU & TRILLES, 2001).

***Emetha* Schioedte and Meinert, 1883;
Emetha audouini (Milne Edwards, 1840)**

Material:

Emetha audouini was collected from the mouth of four of twenty-two *Spicara maena*

(prevalence = 18.2%) on the Erdek coast (Sea of Marmara).

Distribution:

Widely distributed in the Mediterranean and Adriatic Seas (TRILLES, 1994).

Hosts:

Emetha audouini is mainly a parasite of Sparidae and Centracanthidae, particularly *Boops salpa*, *B. boops*, *Spicara vulgaris*, *S. maena*, *S. aprilis*, *S. smaris*, *S. alcedo*, and *S. maurii*, (TRILLES, 1994). Recently reported in cultured *Dicentrarchus labrax* in Greece (PAPAPANAGIOTOU *et al.*, 1999).

***Livoneca* Leach, 1818; *Livoneca punctata* (Uljanin, 1872)**

Material:

Livoneca punctata was collected from the branchial cavity of ten *Alosa fallax* of fifty shads (prevalence = 20%) from the coast of Sinop (Black Sea coast of Turkey) and the Bosphorus (Sea of Marmara). *L. punctata* had previously been recorded in the Black Sea, but this is the first time it is reported amongst Turkish fauna.

Distribution:

TRILLES (1994) lists the known locations for this species: in the Black Sea, particularly the Crimea, the regions of Kerch in Crimea and Gelendzhik, county of Krasnodar in Russia, and region of Agigea, near Constanta, Romania; also reported from Castiglione in the Mediterranean Sea.

Hosts:

Several fishes, particularly Clupeidae and shads: *Clupea pilchardus*, *Alosa finta*, *Caspialosa pontica*, *C. nordmanni*, *Sardinella pilchardus*, *Sardina pilchardus*, *Engraulis encrasicholus ponticus*, *Sprattus sprattus phalericus*, *Trachurus mediterraneus ponticus*; also

Temnodon saltator, *Scorpaena porcus*, *Atherina hepsetus*, *Gobius sp.* (TRILLES, 1994).

***Mothocyta* Costa, in Hope, 1851;**
***Mothocyta epimerica* Costa, 1851**

Material:

Mothocyta epimerica was collected in the branchial cavity of fifteen *Atherina boyeri* of fifty-six specimens (prevalence = 26.8%) from the coast of Sinop (Black Sea of Turkey).

Distribution:

Mothocyta epimerica has been reported in the Mediterranean, Black, and Adriatic Seas (BRUCE, 1986; TRILLES, 1994). Also, recorded from Arcachon, Atlantic Ocean (TRILLES, 1968, 1977).

Hosts:

Exclusively on Atherinidae: *Atherina mochon*, *A. hepsetus*, *A. rissoii* and *A. boyeri* (TRILLES, 1994).

***Mothocyta belonae* Bruce, 1986**

Material:

Mothocyta belonae was collected in the branchial cavity of two *Belone belone* of forty-two specimens (prevalence = 4.8%) from the coast of Sinop (Black Sea coast of Turkey). This species is recorded for the first time in Turkish fauna.

Distribution:

Italy, Livorno. According to BRUCE (1986), "The range of this species within the Mediterranean is not even approximately known".

Hosts:

Recorded only on *Belone belone* (BRUCE, 1986).

List of Cymothoidae parasites and their distribution along Turkish coasts

Cymothoid	Parasitized fish	Location	Author
<i>Anilocra physodes</i> (L. 1758)	Not identified	Mediterranean	MONOD, 1931
	<i>Spicara smaris</i>	Marmara	DEMIR, 1952
	<i>Pagellus</i> sp.	Aegean	GELDIAY & KOCATAS, 1972
	<i>Spicara smaris</i>	Aegean	GELDIAY & KOCATAS, 1972
	Not identified	Mediterranean	TRILLES, 1977
	<i>Scomber japonicus</i>	Aegean	AKMIRZA, 1997
	<i>Sparus auratus</i>	Aegean	KIRKIM, 1998
	<i>Dicentrarchus labrax</i>	Aegean	KIRKIM, 1998
	<i>Boops boops</i>	Aegean	AKMIRZA, 2000
	<i>Diplodus annularis</i>	Aegean	AKMIRZA, 2000
	<i>Diplodus vulgaris</i>	Aegean	AKMIRZA, 2000
	<i>Diplodus sargus</i>	Aegean	AKMIRZA, 2000
	<i>Pagellus erythrinus</i>	Aegean	AKMIRZA, 2000
	<i>Spondyliosoma cantharus</i>	Aegean	AKMIRZA, 2000
	<i>Oblada melanura</i>	Aegean	AKMIRZA, 2000
	<i>Spicara maena</i>	Aegean	AKMIRZA, 2001
<i>Anilocra</i> sp.	<i>Dicentrarchus labrax</i>	Aegean	YELER, 1998
<i>Anilocra frontalis</i> Milne Edwards, 1840	<i>Syphodus tinca</i>	Black	Present study
<i>Nerocila bivittata</i> (Risso, 1816)	<i>Pagellus erythrinus</i>	Mediterranean	MONOD, 1931
	<i>Pagellus</i> sp.	Marmara	DEMIR, 1952
	<i>Pagellus</i> sp.	Aegean	GELDIAY & KOCATAS, 1972
	<i>Sparus auratus</i>	Aegean	KIRKIM, 1998
	<i>Gobius niger</i>	Aegean	KIRKIM, 1998
	<i>Sciaena umbra</i>	Aegean	KIRKIM, 1998
<i>Nerocila maculata</i> (Milne Edwards, 1840)	Not identified	Mediterranean	TRILLES, 1977
<i>Nerocila orbignyi</i> (Guérin-Meneville, 1828-1832)	<i>Dicentrarchus labrax</i>	Aegean	HORTON & OKAMURA, 2001
	<i>Liza aurata</i>	Black	Present study
<i>Ceratothoa oestroides</i> (Risso, 1826)	Not identified	Mediterranean	TRILLES, 1977
	<i>Scomber japonicus</i>	Aegean	AKMIRZA, 1997
	<i>Spicara maena</i>	Aegean	KIRKIM, 1998
	<i>Boops boops</i>	Aegean	KIRKIM, 1998
	<i>Diplodus annularis</i>	Aegean	KIRKIM, 1998
	<i>Sparus auratus</i>	Aegean	KIRKIM, 1998
	<i>Raja alba</i>	Aegean	KIRKIM, 1998
	<i>Trachurus mediterraneus</i>	Aegean	AKMIRZA, 1998
	<i>Boops boops</i>	Aegean	AKMIRZA, 1998

List of Cymothoidae cont'd

Cymothoid	Parasitized fish	Location	Author
	<i>Sparus auratus</i>	Aegean	TOKSEN, 1999
	<i>Dicentrarchus labrax</i>	Aegean	TOKSEN, 1999
	<i>Mullus surmuletus</i>	Aegean	AKMIRZA, 2000
	<i>Diplodus annularis</i>	Aegean	AKMIRZA, 2000
	<i>Diplodus vulgaris</i>	Aegean	AKMIRZA, 2000
	<i>Spicara smaris</i>	Aegean	AKMIRZA, 2001
	<i>Dicentrarchus labrax</i>	Aegean	HORTON & OKAMURA, 2001, 2003
	<i>Spicara maena</i>	Marmara, Bosphorus	Present study
	<i>Sardina pilchardus</i>	Marmara	Present study
<i>Ceratothoa parallela</i> (Otto, 1828)	<i>Boops boops</i>	Aegean	GELDIAY & KOCATAS, 1972
	<i>Boops boops</i>	Aegean	AKMIRZA, 1998
	<i>Citharus macrolepidotus</i>	Aegean	KIRKIM, 1998
	<i>Diplodus annularis</i>	Aegean	KIRKIM, 1998
	<i>Spicara maena</i>	Aegean	KIRKIM, 1998
	<i>Boops boops</i>	Aegean	KIRKIM, 1998
	<i>Boops boops</i>	Marmara	Present study
<i>Ceratothoa italicica</i> Schioedte & Meinert, 1883	<i>Dicentrarchus labrax</i>	Aegean	Present study
	<i>Spicara maena</i>	Bosphorus	Present study
<i>Ceratothoa capri</i> (Trilles, 1964)	<i>Diplodus annularis</i>	Aegean	KIRKIM, 1998
	<i>Boops boops</i>	Aegean	KIRKIM, 1998
	<i>Sparus auratus</i>	Aegean	KIRKIM, 1998
	<i>Puntazzo puntazzo</i>	Aegean	KIRKIM, 1998
<i>Ceratothoa</i> sp.	<i>Diplodus annularis</i>	Aegean	AKMIRZA, 2000
	<i>Oblada melanura</i>	Aegean	AKMIRZA, 2000
	<i>Boops boops</i>	Aegean	AKMIRZA, 2000
	<i>Sarpa salpa</i>	Aegean	AKMIRZA, 2000
	<i>Mullus surmuletus</i>	Aegean	AKMIRZA, 2000
	<i>Scomber japonicus</i>	Aegean	AKMIRZA, 2001
	<i>Spicara smaris</i>	Aegean	AKMIRZA, 2001
<i>Emetha audouini</i> (Milne Edwards, 1840)	<i>Boops boops</i>	Aegean	KIRKIM, 1998

List of Cymothoidae cont'd

Cymothoid	Parasitized fish	Location	Author
	<i>Spicara smaris</i>	Aegean	KIRKIM, 1998
<i>Livoneca punctata</i> (Uljanin, 1872)	<i>Alosa fallax</i>	Black, Bosphorus	Present study
<i>Mothocya epimerica</i> Costa, 1851	<i>Atherina boyeri</i>	Black	ÖKTENER & SEZGIN, 2000
	<i>Atherina boyeri</i>	Black	Present study
<i>Mothocya belonae</i> Bruce, 1986	<i>Belone belone</i>	Black	Present study

List of parasitized fish with corresponding species of Cymothoidae

Parasitized fish	Cymothoidae	Parasitized fish	Cymothoidae
<i>Alosa fallax</i>	<i>Livoneca punctata</i>	<i>Pagellus erythrinus</i>	<i>Anilocra physodes</i>
<i>Atherina boyeri</i>	<i>Mothocya epimerica</i>	<i>Pagellus</i> sp.	<i>Anilocra physodes</i>
<i>Belone belone</i>	<i>Mothocya belonae</i>		<i>Nerocila bivittata</i>
<i>Boops boops</i>	<i>Anilocra physodes</i>	<i>Puntazzo puntazzo</i>	<i>Ceratothoa capri</i>
	<i>Ceratothoa oestroides</i>	<i>Raja alba</i>	<i>Ceratothoa oestroides</i>
	<i>Ceratothoa parallelia</i>	<i>Sardina pilchardus</i>	<i>Ceratothoa oestroides</i>
	<i>Ceratothoa capri</i>	<i>Sarpa salpa</i>	<i>Ceratothoa</i> sp.
	<i>Ceratothoa</i> sp.	<i>Sciaena umbra</i>	<i>Nerocila bivittata</i>
	<i>Emetha audouini</i>	<i>Scomber japonicus</i>	<i>Anilocra physodes</i>
<i>Citharus macrolepidotus</i>	<i>Ceratothoa parallelia</i>		<i>Ceratothoa oestroides</i>
<i>Dicentrarchus labrax</i>	<i>Anilocra physodes</i>		<i>Ceratothoa</i> sp.
	<i>Anilocra</i> sp.	<i>Sparus auratus</i>	<i>Anilocra physodes</i>
	<i>Ceratothoa italicica</i>		<i>Ceratothoa oestroides</i>
	<i>Ceratothoa oestroides</i>		<i>Ceratothoa capri</i>
	<i>Nerocila orbignyi</i>		<i>Nerocila bivittata</i>
<i>Diplodus annularis</i>	<i>Anilocra physodes</i>	<i>Spicara maena</i>	<i>Anilocra physodes</i>
	<i>Ceratothoa oestroides</i>		<i>Ceratothoa italicica</i>
	<i>Ceratothoa parallelia</i>		<i>Ceratothoa oestroides</i>
	<i>Ceratothoa capri</i>		<i>Ceratothoa parallelia</i>
	<i>Ceratothoa</i> sp.		<i>Emetha audouini</i>
<i>Diplodus sargus</i>	<i>Anilocra physodes</i>	<i>Spicara smaris</i>	<i>Anilocra physodes</i>
<i>Diplodus vulgaris</i>	<i>Anilocra physodes</i>		<i>Ceratothoa oestroides</i>
	<i>Ceratothoa oestroides</i>		<i>Ceratothoa</i> sp.
	<i>Nerocila bivittata</i>		<i>Emetha audouini</i>
<i>Gobius niger</i>	<i>Nerocila orbignyi</i>	<i>Spondylisoma cantharus</i>	<i>Anilocra physodes</i>
<i>Liza aurata</i>	<i>Ceratothoa oestroides</i>	<i>Syphodus tinca</i>	<i>Anilocra frontalis</i>
<i>Mullus surmuletus</i>	<i>Ceratothoa</i> sp.	<i>Trachurus mediterraneus</i>	<i>Ceratothoa oestroides</i>
<i>Oblada melanura</i>	<i>Anilocra physodes</i>		
	<i>Ceratothoa</i> sp.		

ACKNOWLEDGEMENTS

We thank the reviewers for their time to carefully review our manuscript. We believe that

their positive comments substantially improved this article.

REFERENCES

- AKMIRZA, A. 1997. The parasites of chub mackerel (*Scomber japonicus*). J. Ege Univ. Aquatic Sci., 14(1-2): 173-181.
- AKMIRZA, A. 1998a. The parasite fauna of horse mackerel. III. National Symposium of Aquatic Products, Ziraat Fakültesi Su Ürünleri Bölümü, Erzurum, 10-12 June 1998, pp. 333-344.
- AKMIRZA, A. 1998b. Parasites in bogue (*Boops boops* L., 1758). J. Ege Univ. Aquatic Sci., 15(3-4): 183-198.
- AKMIRZA, A. 2000a. Metazoon parasites of red mullet (*Mullus surmuletus*) caught near Gökçeada. J. St. Univ. Vet. Fac., 26(1): 129-140.
- AKMIRZA, A. 2000b. Seasonal distribution of parasites detected in fish belonging to the Sparidae family found near Gökçeada. J. Parasitol. Turkey, 24(1): 435-441.
- AKMIRZA, A. 2001. The samples from metazoon parasites detected in fish around Gökçeada. Congress of National Aegean Islands, 7, pp. 85-96.
- BORCEA, I. 1933. *Livoneca pontica*, nov. sp., cymothoide parasite des aloses et sardines de la Mer Noire. Ann. Sci. Univ. Jassy, XVII(3-4): 481-502
- BRUCE, N.L. 1986. Revision of the Isopod crustacean genus *Mothocya* Costa, in Hope, 1851 (Cymothoidae: Flabellifera), parasitic on marine fishes. J. Natl. Hist., 20: 1089-1112.
- BRUCE, N.L. 1987. Australian species of *Nerocila* Leach, 1818 and *Creniola* n.gen. (Isopoda: Cymothoidae), crustacean parasites of marine fishes. Records of the Australian Museum, 39: 355-412.
- BRUSCA, R.C. 1981. A monograph of the Isopode Cymothoidae (Crustacea) of the eastern Pacific. Zool. Journal of the Linnean Society, 73: 117-199.
- COLORNI, A., J.-P. TRILLES & D. GOLANI. 1997. *Livoneca* sp. (Flabellifera: Cymothoidae), an isopod parasite in the oral and branchial cavities of the Red Sea silverside *Atherinomorus lacunosus* (Perciformes, Atherinidae). Dis. Aquat. Org., 31: 65-71.
- DEMIR, M. 1952. The benthic Invertebrates of the Bosphorus and Islands Coasts. J. St. Univ. Fen Fak. Hidrobiol. A: 362-363.
- GELDIAY, R. & A. KOCATAS. 1972. Isopods collected in Izmir Bay, Aegean Sea. Crustaceana, Suppl. 3, Studies on Peracarida: 19-30.
- HOLLAND, C.V. & C.R. KENNEDY. 1997. A checklist of parasitic helminth and crustacean species recorded in freshwater fish from Ireland. Biology and Environment: Proceedings of the Royal Irish Academy, 97B(3): 225-243.
- HORTON, T. 2000. *Ceratothoa steindachneri* (Isopoda: Cymothoidae) new to British waters with a key to north-east Atlantic and Mediterranean *Ceratothoa*. J. Mar. Biol. Ass. U.K., 80: 1041-1052.
- HORTON, T. & B. OKAMURA. 2001. Cymothoid isopod parasites in aquaculture: a review and case study of a Turkish sea bass (*Dicentrarchus labrax*) and sea bream (*Sparus auratus*) farm. Dis. Aquat. Org., 46: 181-188.
- HORTON, T. & B. OKAMURA. 2003. Post-haemorragic anaemia in sea-bass, *Dicentrarchus labrax* (L.), caused by blood feeding of *Ceratothoa oestroides* (Isopoda: Cymothoidae). J. Fish Dis., 26: 401-406.
- KIRKIM, F. 1998. Investigations on Systematic and Ecology of Isopoda Fauna in the Aegean Sea. D.E.U. Thesis. Marine Science and Technology Institution, Izmir, 80 pp.
- MLADINEO, I. 2002. Prevalence of *Ceratothoa oestroides* (Risso, 1826), a cymothoid isopode parasite, in cultured sea bass *Dicentrarchus*

- labrax* L. on two farms in middle Adriatic Sea. Acta Adriat., 43(1): 97 -102
- MONOD, T. 1931. Crustacés de Syrie. In: A. Gruvel. Les Etats de Syrie. Richesses marines et fluviales. Exploitation actuelle. Avenir. Bibl. Faune Colon. Fr., 397-435.
- MOREIRA, P. S. & V. SADOWSKY. 1978. An annotated bibliography of parasitic Isopoda (Crustacea) of Chondrichthyes. Bolm. Inst. Oceanogr., Sao Paulo, 27(2): 95-152.
- ÖKTENER A. & M. SEZGIN. 2000. *Mothocya epimerica* Costa, 1851 (Flabellifera: Cymothoidae), an isopod parasite in the branchial cavities of the Black Sea Silverfish *Atherina boyeri* Risso, 1810 (Perciformes, Atherinidae). Turkish J. Marine Sciences, 6 (1): 23-29.
- PAPAPANAGIOTOU, E.P. & J.-P. TRILLES. 2001. Cymothoid parasite *Ceratothoa parallela* inflicts great losses on cultured gilthead seabream *Sparus aurata* in Greece. Dis. Aquat. Organisms, 45: 237-239.
- PAPAPANAGIOTOU, E.P., J.-P. TRILLES & G. PHOTIS. 1999. First record of *Emetha audouini*, a cymothopid isopod parasite from cultured sea bass *Dicentrarchus labrax* in Greece. Dis. Aquat. Org., 38: 235-237.
- PAPOUTSOGLOU S., M.J. COSTELLO, E. STAMOU & G. TZIHA. 1996. Environmental conditions at sea cages, and ectoparasites on farmed European sea bass *Dicentrarchus labrax* and gilt-head sea bream *Sparus aurata* at two farms in Greece. Aquacult. Res., 27: 25-34.
- ROMESTAND, B. 1978. Etude Écophysiologique des Parasitoses à Cymothoidae. Thesis, Université de Montpellier, France, 284 pp.
- ROMESTAND, B. 1979. Etude Écophysiologique des Parasitoses à Cymothoadien. Ann. Parasitol. Hum. Comp., 54(4): 423-448.
- ROMESTAND, B. & J.-P. TRILLES. 1976. Production d'une substance anticoagulante par les glandes endocrines céphalothoraciques des Isopodes Cymothoidae *Meinertia oestroides* (Risso, 1826) et *Anilocra physodes* (L. 1758) (Isopoda, Flabellifera, Cymothoidae). C. R. Hebd. Séanc. Acad. Sci., Paris, 282: 663-665.
- ROMESTAND, B. & J.-P. TRILLES. 1977. Influence des cymothoadiens (Crustacea, Isopoda, Flabellifera) sur certaines constantes hématoLOGIQUES des poissons hôtes. Zeitschrift für Parasitenkunde, 52: 91-95.
- ŠARUŠIĆ, G. 1999. Preliminary report of infestation by isopod *Ceratothoa oestroides* (Risso, 1826), in marine cultured fish. Bull. Eur. Assoc. Fish Pathologists, 19: 110-112.
- SIRIKANCHANA, P. 1982. Check lists of parasites of fishes in Thailand. Notes from the Faculty Fisheries, Kasetsart University, Bangkok, 10: 1-11.
- TOKSEN, E. 1999. Metazoon Gill Parasites of Culture Gilthead Sea Bream (*Sparus aurata* L.) and Sea Bass (*Dicentrarchus labrax* L.) in Aegean Sea Coast and their Treatment. Thesis, Ege Univ. Sci. Inst., Izmir, 153 pp.
- TRILLES, J.-P. 1965. Sur deux espèces d'Anilocres (Isopodes, Cymothoidae) mal connues: *Anilocra physodes* L. et *Anilocra frontalis* (Milne Edwards). Ann. Parasit. Hum. Comp., 40(5): 575-594.
- TRILLES, J.-P. 1968. Recherches sur les Isopodes Cymothoidae des côtes Françaises. Thesis, Univ. Montpellier, France, N° enregistrement CNRS AO 2305. Vol. 1: Bionomie et parasitisme, Vol. 2: Biologie générale et sexualité, 793 pp. Deuxième thèse, Systématique et Faunistique, 181 pp.
- TRILLES, J.-P. 1969. Recherches sur les Isopodes Cymothoidae des côtes Françaises. Aperçu général et comparatif sur la bionomie et la sexualité de ces crustacés. Bull. Soc. Zool. Fr., 94(3): 433-445.
- TRILLES, J.-P. 1972. Les Cymothoidae (Isopoda, Flabellifera) des côtes Françaises (systématique, faunistique, écologie et répartition géographique). I. Les Ceratothoinae Schioedte et Meinert, 1883. Bull. Mus. Natl. Hist. Nat., Paris, 3ème série, N° 91 (Zool. 70): 1191-1230.
- TRILLES, J.-P. 1975. Les Cymothoidae (Isopoda, Flabellifera) des collections du Muséum National d'Histoire Naturelle de Paris. II. Les Anilocridae Schioedte et Meinert, 1881. Genre *Anilocra*, Leach, 1818 et *Nerocila*, Leach, 1818. Bull. Mus. Natl. Hist. Nat., Paris, 3ème série, 290 (Zool. 200): 303-346.

- TRILLES, J.-P. 1976. Les Cymothoidae (Isopoda, Flabellifera) des côtes Françaises. III. Les Lironecinae Schioedte et Meinert, 1884. Bull. Mus. Natl. Hist. Nat., Paris, 3ème série, 390 (Zool. 272): 977-993.
- TRILLES, J.-P. 1977. Les Cymothoidae (Isopoda, Flabellifera) parasites des poissons du Rijksmuseum van Natuurlijke Historie de Leiden. Méditerranée et Atlantique Nord-Oriental. Zoologische Mededelingen, Leiden, 52(2): 7-17.
- TRILLES, J.-P. 1979. Les Cymothoidae (Isopoda, Flabellifera) parasites de poissons du Rijksmuseum van Natuurlijke Historie de Leiden.
- II. Afrique, Amérique et régions Indo-ouest-Pacifiques. Zoologische Mededelingen, Leiden, 54(17): 245-275.
- TRILLES, J.-P. 1994. Les Cymothoidae (Crustacea, Isopoda) du monde (prodrome pour une faune). Studia Marina, 21/22(1-2): 1-288.
- TRILLES, J.-P., B.M. RADUJKOVIĆ & B. ROMESTAND. 1989. Parasites des poissons marins du Monténégro: Isopodes. Acta Adriat., 30(1/2): 279-306.
- YELER, S. 1998. Investigation of the Parasites of Sea Bass (*Dicentrarchus labrax* L.) under Culture Conditions in Bodrum. Master Thesis, Ank. Univ. Sci. Inst., 46 pp.

Received: 11 November 2003

Accepted: 28 September 2004

Popis cimotoidnih rakova (Crustacea, Isopoda) na morskim ribama Turske

Ahmet ÖKTENER¹ i Jean-Paul TRILLES²

¹ Cihannüma mah. Hüsnü savman sokak no: 22/5 80690 Beşiktaş, İstanbul, Turska

²UMR 5171 (Genom, populacija, interakcija, adaptacija), Ekipa ekofiziološke adaptacije i ontogeneze, Sveučilište u Montpellier-u 2, CC. 092, Place E. Bataillon, 34095 Montpellier Cedex O5, Francuska

SAŽETAK

U radu se daje opis cimotoidnih vrsta rakova u Turskoj. Neke vrste su sabrali autori kao što su *Anilocra frontalis* (Milne Edwards, 1840); *Nerocila orbignyi* (Guérin-Méneville, 1829-1832); *Ceratothoa italicica* (Schioedte i Meinert, 1883); *Ceratothoa oestroides* (Risso, 1826); *Ceratothoa parallela* (Otto, 1828); *Emetha audouini* (Milne Edwards, 1840); *Livoneca punctata* (Uljanin, 1872); *Mothocyta epimerica* (Costa, 1851), i *Mothocyta belonae* (Bruce, 1986). Četiri od spomenutih vrsta po prvi su put zabilježene u fauni Turske i to: *A. frontalis*, *C. italicica*, *L. punctata*, i *M. belonae*. Lista cimotoidnih rakova, nađenih na ribama Turske, zajedno s novim nalazima, obuhvaća 13 vrsta (2 još nedeterminirane) iz 6 rodova (*Anilocra*, *Nerocila*, *Ceratothoa*, *Emetha*, *Livoneca*, i *Mothocyta*), i 27 ribljih domaćina. Izneseni su podaci o distribuciji i ribljim domaćinima, te odgovarajući autori.

Ključne riječi: Cymothoidae, Crustacea, parazitski izopodi, Turska