

---

---

**Distribution of juvenile anchovy *Engraulis encrasicolus* (L.) in an estuarine habitat and influence of year–class strength on its catch value**

G. Sinovčić

*Institute of Oceanography and Fisheries, Split, Croatia*

---

*Distribution and growth of anchovy, *Engraulis encrasicolus* (L.) and year – class strength appearance were investigated in the Zrmanja river estuary (middle part of the Eastern Adriatic). Influence of anchovy strong age - class on the exploitation of adults was studied as well. Monthly representative samples were taken from the eastern (Croatian) part of the Middle Adriatic (Novigrad Sea) from the initial appearance of strong year classes in August 1989 in other words July 1990 up to March 1991. Catch data obtained during the 1979 – 1993 period were used as well.*

---

INTRODUCTION

Usually, recruitment of fish population is predicted more reliably by abundances of life stages other than eggs and larvae (Sissenwine, 1984; Smith, 1985; Sale, 1990; Bailey, 1994). Year class strength is primarily estimated during the first year of life and depends on the conditions for fish reproduction and survival during the early developmental stages of ontogeny (Hjort, 1914; Deckhnik *et al.*, 1985). Survival during the early stages of pelagic fish is usually attributed to environmental factors, but may in part be also due to fluctuations in the abundances of particular age classes of predators on the eggs, larvae and

juveniles of either the parent stock or some other species. Therefore potential causes of variations in recruitment to fish stocks are:

- variations in environmental factors including climate, tidal conditions, etc.
- variations in the abundance of the parent stock as a whole (the stock-recruitment effect)
- variations in the abundance of predators on, or competitors of the pre-recruit stages (ecological interactions)

The perceived abundance of the stock will be influenced primarily by the variations of recruitment, compounded by the effects of fishing and possible shifts of geographical range. It is difficult to predict year-class strength until the weather that affects it has actually happened. This would therefore give us an advantage of most a year of two on use a 0- or 1- group survey.

Anchovy landings have decreased greatly since 1985 without a considerable change in the fishing effort (Sinovčić *et al.*, 1991, Sinovčić, 1992). Attempts have been made to explain these fluctuation in a variety of ways (Regner *et al.*, 1981; Sinovčić *et al.*, 1991; Sinovčić, 1992; Sinovčić and Alegria-Hernandez, 1997).

The main objectives of this study were, first, to determine the year-class strength of juvenile anchovy in the estuarine habitat (Zrmanja river estuary), its distribution, and second, to identify its effect on the catch of adults in forthcoming years.

#### Area characteristics

The values of phosphate which totalled  $0.25 \mu\text{mol l}^{-1}$  and especially silicates which amounted to  $12.8 \mu\text{mol l}^{-1}$  in the Novigrad Sea and  $30.8 \mu\text{mol l}^{-1}$  in the Karin Sea (Table 1) make this area one of the richest in the eastern part of the Adriatic, which, is generally speaking, oligotrophic due to the quantity of nutrients.

Considering the categorization of some eastern parts of the Adriatic ecosystem on the basis of the phytoplankton cell volume, Viličić (1989) noted the highest values in the Novigrad Sea and the Karin Sea bays as well as, in an urban area of Šibenik bay. Based on the phytoplankton quantity, the Novigrad Sea and the Karin Sea bays are classed among the highest category of naturally eutrophicated areas. In that respect, only the Gruž and Šibenik harbour are richer. However, high production in these areas is the result of urban eutrophication, while the Novigrad Sea and the Karin Sea bays are naturally eutrophicated

Table 1. Physical-chemical characteristics and nutrient concentrations in the Karin and the Novigrad Sea (Buljan, 1969) as well as, in the Velebit Channel and the Nin Bay (Škrivanić and Barić, 1979)

Area	S x 10 <sup>-3</sup>	Temp. <sup>o</sup> C Min-max	PO <sub>4</sub> -P μmol l <sup>-1</sup>	NO <sub>3</sub> -N μmol l <sup>-1</sup>	SiO <sub>2</sub> -Si μmol l <sup>-1</sup>
Karin Sea	19.52-36.60	4.1-23.4	0.00-0.25	-	3.89-30.80
Novigrad Sea	11.72-36.92	4.6-23.0	0.00-0.25	-	10.00-12.80
Velebit Channel	34.10-37.92	8.0-23.1	0.02-0.05	0.02-0.60	0.60-4.00
Nin Bay	36.50-38.10	7.9-24.7	0.02-0.06	0.14-0.40	1.00-4.00

## MATERIAL AND METHODS

During the periods of study, 2564 juvenile anchovy from the interconnected bays of the Novigrad Sea and Karin Sea (Fig. 1) were collected.

Monthly representative anchovy samples were obtained by purse seine catches from August 1989 to March 1991.

The total length of each fish was determined, to a mm below its actual length. The gonads and gut content were then removed and the fish weighed to the nearest 0.1 g. A total of analyzed juvenile anchovy specimens ranged from 4.4 to 12.5 cm and from 0.4 to 13.8 g.

The otoliths were extracted, washed and placed in denoted envelopes.

For studies of the anchovy state under exploitation, fishermen were requested to fill up the forms recording daily catch by fishing ground in the Novigrad Sea.

Total catch data of the eastern and western Adriatic were available from FAO yearbooks of fishery statistics.

## RESULTS AND DISCUSSION

### Distribution of juvenile anchovy

Juvenile anchovy started to appear in mass (year – class strength) and to be fished along the eastern Adriatic in July 1989 and August 1990. The species of small pelagics react very quickly to the increase of

phytoplankton and zooplankton production. Due to their position in the trophic chain, they influence the trophic relations in the sense that they carry organic matter from lower to higher trophic levels.

All analyzed juvenile anchovy specimens ranged from 4.4 to 12.5 cm and from 0.4 to 13.8 g. They are in first year of life.

Juvenile anchovies remain in the Novigrad Sea throughout the year using it as a nursery area. Juveniles remain in this area before migrating into deeper waters as they increase in size. Data of realized juvenile anchovy catches which are represented on Fig. 1. simultaneously show their spatial – temporal distribution in this area.




Fig. 1. Map showing the investigation area as well as the spatial and temporal distribution of juvenile anchovy in the Novigrad Sea during 1990

In spring, schools of this species were located near the entrance to Novigrad Sea.

During the summer, they extended their presence to the south and the west coasts of the bay, which are better afforested and hence, the more productive coasts of the bay. Organic matter accompanies the rain,

enriching the area with nutrients which fulfil the basic precondition of phytoplankton, as well as zooplankton swelling. The south and west coasts are the most afforested. They are the richest and most productive parts of the area. Results of the tintinnines quantity, which comprise the food of the juvenile anchovy (Ercegović, 1940), showed increased amounts in the Novigrad Sea, particularly during spring and summer, when compared with other parts of the eastern Adriatic (Kršinić, 1987). The amounts show that these groups of organisms make this aquatorium one of the richest areas, which is very important due to the fact that plankton organisms provide basic food for pelagic species.

In autumn they migrate to deeper waters; but in winter they move to the deepest, central part of the bay, where they find the most favorable temperature conditions.

### Catches

Figure 2. shows how the strong age-class which appeared in 1989




Fig. 2. Fluctuations of annual anchovy catches in the Adriatic during 1979 – 1993 period

and 1990 and observed during constant monitoring, influenced the overall catches of this species in the eastern part of the Adriatic, since these fish later become (under normal conditions) the bulk of the Adriatic anchovy stock. It is therefore expected that this year – class strength of anchovy recorded in 1989 and 1990 will significantly reflect upon increased catches during subsequent fishing seasons depending on anchovy life – span. The catches realized in the western part of the Adriatic as well as catches for the entire Adriatic Sea are shown as well. Since 1989 and 1990, with results of the existence of strong year classes, anchovy catches have been increased. As the anchovy life span is five years (Sinovčić, 1988, 1991, 1992), the year – class strength appearance has influenced the catch value during the period 1990 to 1993. (Fig. 2).

## CONCLUSIONS

The results of this study showed the following: Novigrad Sea and Karin Sea bays constitute a favourable nursing habitat for juvenile anchovy.

Juvenile anchovy have a coastal distribution during and summer. During autumn they migrate to deeper waters, but in winter they move to the deepest, central part of the bay, where they find the most favourable temperature conditions.

The importance of the influence of anchovy year – class strength and its successive catches depending on the anchovy life span was asserted.

## REFERENCES

- B a i l e y, K.M. 1994. Predation of juvenile flatfish and recruitment variability. *Netherlands Journal of Sea Research*, 32: 175-189.
- B u l j a n, M, 1969. Neka hidrografska svojstva područja rijeke Krke i Zrmanje. *Krš Jugoslavije*, 6: 303-331.
- D e c k h n i k, T.V., V.P. S e r e b r y a k o v and S.G. S o i n, 1985. Importance of early stages of fish development in the year class abundance formation. In: *Teoriya formirovaniya chislennosti i ratsionalnoga ispolzovaniya promyslovikh ryb*. Nauka press, pp. 56-72. (In Russian).

Ercegović, A. 1940. Ishrana srdele (*Clupea pilchardus* WALB.) u stadiju metamorfoze. God. Oceanogr. Inst., 2: 26-44.

Hjort, J. 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research. Rapp. P-v. Réun. Cons. int. Explor. Mer, 5 (20): 1-228.

Kršinić, F. 1987. Tintinnines (Ciliophora, Oligotrichida, Tintinnina) in Eastern Adriatic Bays. Estuarine, Coastal and Shelf Science, 24: 527-538.

Regner S., C. Piccinetti., M. Specchi and G. Sinovčić. 1981. Preliminary statistical analysis of sardine stock estimation from data obtained by egg surveys. FAO Fish. Rep., 253: 143-154.

Sale, P. 1990. Recruitment in marine species: in the bandwagon rolling in the right direction? Trends in Ecology and Evolution, 5: 25-27.

Sinovčić, G. 1988. Age and growth of anchovy, *Engraulis encrasicolus* (L.) in the Middle Adriatic. Rapp. Comm. Int. Mer Médit., 31 (2): p. 266.

Sinovčić, G. 1992. Biologija i dinamika populacije brgljuna. *Engraulis encrasicolus* (Linnaeus, 1758) u Jadranu. PhD Thesis, University of Zagreb, 163 pp. (mimeo)

Sinovčić, G., V. Alegria and I. Jardaš. 1991. Biološka osnova pelagijskog i priobalnog ribolova Jugoslavije. Pomorski zbornik, 29: 383-408.

Sinovčić, G., V. Alegria – Hernandez. 1997. Variation in abundance and size of the sardine, *Sardina pilchardus* (Walbaum) in the eastern Adriatic. Oceanologica acta, 20 (1): 201-206.

Sissenwine, M.P. 1984. Why do fish population vary? In: Exploitation of Marine Communities. MAY, R.M., (Editor). Springer Verlag, New York, N.Y., 59-94.

S m i t h, P.E. 1985. Year class strength and survival of 0-group clupeoids. Canadian Journal of Fisheries and Aquatic Sciences, 42 (Suppl. 1): 69-82.

Š k r i v a n i ć, A. and A. B a r i ć. 1979. Cruises of the research vessel "Vila Velebita" in the Kvarner of the Adriatic Sea. IV. Distribution of the primary nutrients. Thalassia Jugosl., 15: 61-88.

V i l i č i ć, D. 1989. Phytoplankton population density and volume as indicators of eutrophication in the eastern part of the Adriatic Sea. Hydrobiologia, 174: 117-132.

Accepted: 15 April 1998.



# Rasprostranjenost juvenilna brgljuna *Engraulis encrasicolus* (L.) u estuariju i utjecaj pojave snažne dobne skupine na njegovo iskorištavanje

G. Sinovčić

*Institut za oceanografiju i ribarstvo, Split, Hrvatska*

## SAŽETAK

U radu se iznose rezultati istraživanja rasprostranjenosti juvenilna brgljuna, *Engraulis encrasicolus* (L.) u estuariju rijeke Zrmanje (Novigradsko i Karinsko more – središnji dio istočnoga dijela Jadrana). Prati se utjecaj snažne dobne skupine brgljuna, od njezine prve pojave (kolovoz 1989) do prelaska juvenilna brgljuna u adultnu fazu (ožujak 1990, odnosno 1991), na ukupni ulov ove značajne pelagičke vrste.

*Engraulis encrasicolus* (L.) u estuariju  
utjecaj pojave snažne dobnе stępnе na  
njegovu inkubacijsku i razvojnu dinamiku  
U radu se istražuje utjecaj različitih faktora na inkubacijsku i razvojnu dinamiku  
ikre i mlade ribice *Engraulis encrasicolus* (L.) u estuariju rijeke Zravnice  
(Zravnica, Kvarneri zaton - estuarij rijeke Zravnice).  
Istraženo je dobnе stępnе i vrijeme inkubacije, od kojih je najveći utjecaj  
imao na inkubacijsku i razvojnu dinamiku. U radu se istražuje utjecaj  
različitih faktora na inkubacijsku i razvojnu dinamiku ikre i mlade ribice  
1990. godine (1991. na otvorenom moru) u estuariju rijeke Zravnice.  
U radu se istražuje utjecaj različitih faktora na inkubacijsku i razvojnu  
dinamiku ikre i mlade ribice u estuariju rijeke Zravnice (Zravnica, Kvarneri  
zaton). Istraženo je dobnе stępnе i vrijeme inkubacije, od kojih je najveći  
utjecaj imao na inkubacijsku i razvojnu dinamiku. U radu se istražuje  
utjecaj različitih faktora na inkubacijsku i razvojnu dinamiku ikre i mlade  
ribice 1990. godine (1991. na otvorenom moru) u estuariju rijeke Zravnice.

### Area characteristics

The values of phosphate which totalled 0.25  $\mu\text{mol l}^{-1}$  and especially silicates which amounted to 12.5  $\mu\text{mol l}^{-1}$  in the Rovinj Sea and 30.5  $\mu\text{mol l}^{-1}$  in the Kvarneri Sea (Table 1) make this area one of the richest in the eastern part of the Adriatic, which is generally speaking, oligotrophic due to the scarcity of nutrients. Considering its categorization as being eastern parts of the Adriatic

BILJEŠKE – NOTES, izdaje Institut za oceanografiju i ribarstvo, 21000 Split, Hrvatska;  
Izlazi povremeno.  
Published by Institute of Oceanography and Fisheries, 21000 Split, Croatia;  
Issued periodically.

Glavni urednik:  
I. Marasović  
Editor – in chief: