No. 80

Confirmation of the presence of Gadella maraldi (RISSO, 1810) in the seas of Turkey

Hakan KABASAKAL

University of Istanbul, Faculty of Aquatic Products, Department of Marine Biology, Ordu cad., No: 200, Laleli 34470 Istanbul, Turkey

On November 28, 1997, two female specimens of Gadella maraldi (RISSO, 1810) were caught at a depth of 150 m in the Gulf of Saros (40° 28' 58" N; 26° 10' 30" E). Measurements and meristic counts of this species are given for the first time from Turkish seas. The present recording confirms the presence of G. maraldi in the seas of Turkey.

Key words: Gadella maraldi, Moridae, deep-sea fishes, seas of Turkey

INTRODUCTION

Little is known about the deep-sea fishes in the seas of Turkey and very few studies have been carried out on this subject (DEMİR, 1958; MATER *et al.*, 1988; KAYA, 1993; ÜNSAL and KABASAKAL, 1998). On November 28, 1997, a bottom trawler haul at 150 m on a muddy-sandy bottom in the Gulf of Saros (40° 28' 58" N; 26° 10' 30" E) (Fig. 1) included two specimens of *Gadella maraldi*

(RISSO, 1810) (Fig. 2) the presence of which was not confirmed until now.



Fig. 1. Sampling station in the Gulf of Saros (40° 28' 58" N; 26° 10' 30" E)




Fig. 2. Gadella maraldi (RISSO, 1810), scale bar 10 mm

G. maraldi belongs to family Moridae and generally occurs at the depths between 150 and 700 m (COHEN, 1986; BAUCHOT, 1987; COHEN et al., 1990). The occurrence of this species in the eastern Atlantic Ocean (Madeira, coasts of Morocco and Iberian peninsula) and in the Mediterranean Sea has been well documented (ŠOLJAN, 1948, in the Adriatic Sea as Uraleptus maraldi; BENTUVIA, 1971; COHEN, 1986; COHEN et al., 1990). It's presence in the Aegean Sea has been reported by PAPACONSTANTINOU and TSIMENIDIS (1979).

In the present study, a description of *G. maraldi* based on two specimens caught in the Gulf of Saros is given.

MATERIAL AND METHODS

A commercial trawler with a cod-end mesh opening of 22 mm from knot to knot was used for the sampling. Samples were fixed and stored in 5 percent formalin solution. The total lengths of the fishes were 206 and 176 mm, respectively. The specimens were accurately identified according to the taxonomic key provided by COHEN (1986). The total length (TL) was measured to the nearest 1.0 mm, while morphometric measurements to the nearest 0.1 mm. The material was preserved at the Faculty of Aquatic Products, Marine Biology Laboratory at the University of Istanbul.

RESULTS

Selected measurements and meristic counts of the two specimens are presented on Tables 1 and 2.

Table 1. Selected measurements of two specimens; body proportions are given as a percentage of total length

MEASUREMENTS	mm	%	mm	%
Total length	206	-	176	
Standard length	181	87.8	158	89.7
D1 Base length	16.4	7.9	14.05	7.9
D2 Base length	101.4	49.2	88.45	50.2
Interdorsal space	2.75	1.3	1.85	1.05
Base length of anal fin	118.1	57.3	102.35	58.1
Pectoral length	32.85	15.9	31	17.6
Head length	51.3	24.9	41.2	23.4
Eye diameter	14.4	6.9	9.65	5.4
Preorbital distance	12.9	6.2	11.95	6.7
Postorbital distance	24	11.6	19.6	11.1
Interorbital space	16	7.7	12.6	7.1
Body depth at anal fin origin	44	21.3	38.15	21.6
Depth of caudal peduncle	4.55	2.2	3.35	1.9

Table 2. Meristic counts of two specimens

MERISTIC COUNTS	NUMBER			
the states a maximum size of 30 cm.	TL 206 mm	TL 176 mm		
D1 rays	21111	10		
D2 rays	54	54		
Anal rays	57	57		
Pectoral rays	24	20		
Pelvic rays	7	7		
Scales on the lateral line	104	102		
Gill rakers on the 1st left branchial arch	11	12		

Description

The body is elongate, compressed (body depth at anal fin origin 21.3 to 21.6 % of TL) and tapering posteriorly; head is long and 23.4 to 24.9 % of TL. The snout is rather broad and obtusely rounded, preorbital distance 11.1 to 11.6 % of TL. Chin barbel absent. The upper jaw with two rows of teeth, the outer row with small teeth interspersed with notably large ones; the inner row with small teeth only. No teeth on vomer and palatines. A projection of the swimbladder extends far anteriorly on each side of the fish, and attaches to the rear of the cranium. There are two dorsal fins; interdorsal space is narrow and 1.05 to 1.3 % of TL. The origin of the anal fin nearly at the level of the origin of first dorsal fin; anal fin not indented. The tip of each pectoral fin extends far beyond the origin of anal fin. The body and the head covered with small cycloid scales with exception of chin and lips. There is a light organ is present as a small, scaleless patch on the belly.

The body is dark brown; the belly and the sides below the pectoral fins bluish-black. The mouth is pale. The operculum is covered by numerous dark and small spots.

Biology

According to COHEN (1986), BAUCHOT (1987) and COHEN et al. (1990) G. maraldi attains a maximum size of 30 cm, reaches sexual maturity at the total length of 15 cm and spawns in spring. AKŞIRAY (1987) stated that egg diameter of G. maraldi varied between 0.8 and 1.2 mm. No eggs were observed in the ovaries of the specimens with naked eye, but microscopical observations of ovaries revealed numerous eggs which diameter varied between 0.14 and 0.28 mm.

G. maraldi is known to be a carnivorous fish which primarily feeds on benthic invertebrates, such as worms, shrimps and small crustaceans, and small fish (AKŞIRAY, 1987). The stomach of the specimen of 176 mm TL was empty, while the stomach of the specimen of 206 mm TL contained five otoliths and remains of teleost fish (species unidentified).

DISCUSSION AND CONCLUSIONS

The presence of *G. maraldi* in the seas of Turkey is now confirmed, and information about depth and locality where the specimens of this species were caught, are given for the first time from Turkish waters. According to AKŞIRAY (1987) *G. maraldi* is present in the seas of Turkey with exception of Black sea, but AKŞIRAY has not given any information about depth and locality where this species was caught nor where the study material was available for inspection. In one of the most extensive studies on the deep sea fishes of the Aegean Sea, there is no record of *G. maraldi* from the seas of Turkey (KAYA, 1993). Furthermore, there is no record of *G. maraldi* in the recent study of MATER and MERİÇ (1996) which lists marine fishes from the seas of Turkey found to date. PAPACONSTANTINOU (1990) stated *G. maraldi* was a rare fish throughout the Aegean Sea and his suggestion was confirmed by the single occurrence of two specimens during the three year research period.

REFERENCES

- AKŞIRAY, F. 1987. Türkiye Deniz Balıkları ve Tayin Anahtarı. İ. Ü. Rektörlüğü Yayınları No: 3490. İstanbul, 811 pp.
- BAUCHOT, M. -L. 1987. Poisson osseux. In: W. Fischer, M. Schneider and M. -L. Bauchot (Editors). Fiches FAO d'identification des espèces pour les besoins de la pêche. (Révision 1). Méditerranée et mer Noire. Zone de pêche 37. Volume II. Vertébrés. Publication préparée par la FAO et la Commission des Communautés Européennes (Projet GCP/INT/422/EEC) financée conjointement par ces deux organisations. FAO, Rome, pp. 891-1422.
- BEN-TUVIA, A. 1971. Revised list of the Mediterranean fishes of Israel Israel Journal of Zoology, 20: 1-39.
- COHEN, D. M. 1986. Moridae. In: P. J. P. Whitehead, M. -L. Bauchot, J. -C. Hureau, J. Nielsen and E. Tortonese (Editors). Fishes of the North-eastern Atlantic and the Mediterranean, Vol. 2. UNESCO, Paris, pp. 713-723.
- COHEN, D. M., T. INADA, T. IWAMOTO, and N. SCIALABBA. 1990. Gadiform fishes of the world (Order Gadiformes). An annotated and illustrated catalogue of cods, hakes, grenadiers and other gadiform fishes known to date. FAO Fisheries Synopsis, 125: 10, 442 pp.
- DEMİR, M. 1958. Marmara ve kuzey-doğu Ege'den üç derin deniz balığı nev'i. Hidrobiologi Mecmuası, Seri A, 4: 134-151.
- KAYA, M. 1993. An investigation on the deep sea fishes of the Aegean Sea. Tr. J. of Zoology, 17: 411-426.
- MATER, S., M. KAYA and H. A. BENLİ. 1988. An investigation on the deep sea (bathyal) fishes of the Gökova Bay, Aegean Sea. Rapp. Comm. Int. Mer Médit., 31, p. 276.
- MATER, S. and N. MERİÇ. 1996. Deniz Balıkları (Pisces). In: A. Kence and C. C. Bilgin (Editors). Türkiye Omurgalılar Tür Listesi. TÜBİTAK, Ankara, pp. 133-172.
- PAPACONSTANTINOU, C. and N. TSIMENIDIS. 1979. Some uncommon fishes from the Aegean Sea. Cybium 3e série, 7: 3-14.

- PAPACONSTANTINOU, C. 1990. Some rare mesopelagic and bathyal fishes caught in the Greek Seas. Thalassographica, 13: 35-39.
- ŠOLJAN, T. 1948. Fishes of the Adriatic (Ribe Jadrana). Fauna et Flora Adriatica, 1. Nakladni Zavod Hrvatske. Zagreb, 428 pp.
- ÜNSAL, N. and H. KABASAKAL. 1998. New records for the deep sea fish fauna of Gökçeada. Celal Bayar University. Journal of Science and Arts Faculty, The Series of Natural Sciences (Biology), 1: 59-62.

Accepted: 25 November 1998

Potvrda prisutnosti *Gadella maraldi* (RISSO, 1810) u turskim vodama

Hakan KABASAKAL

Sveučilište u Istanbulu, Fakultet vodenih resursa, Odjel biologije mora, Ordu. cad., No: 200, Laleli 34470 Istanbul, Turska

SAŽETAK

U kočarskoj lovini ostvarenoj 28. studenoga 1997. u Saroskom zaljevu (Egejsko more) na 150 m dubine i muljevito-pjeskovitom dnu ulovljena su dva primjerka ribe *Gadella maraldi* (Risso, 1810) (por. Moridae), što je potvrda njene prisutnosti u turskim vodama. Ukupna tjelesna dužina primjeraka bila je 206, odnosno 176 mm. Iznose se njihove morfometrijske i merističke karakteristike.